



Environment and Time as Constraints on the Biogeographical Distribution of Gibbons

Journal:	<i>American Journal of Primatology</i>
Manuscript ID	AJP-18-0040.R2
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Dunbar, Robin; Psychology Cheyne, Susan; Borneo Nature Foundation Lan, Daoying; Guangdong Institute of Applied Biological Resources Korstjens, Amanda; Bournemouth University, School of Applied Sciences Lehmann, Julia; Roehampton University, Human and Life Sciences Cowlshaw, Guy; Institute of Zoology
Indicate which taxonomic group was the subject of your study (select all that apply or type another option)::	Apes (non-human)
Keywords:	climate, biogeographic distribution, group size, foraging ecology

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Highlights:

- Time budgets limit the biogeographic distribution of gibbons and siamang
- Time that has to be allocated to resting seems to be the main constraint
- The limits on group size are not so restrictive as to make monogamy obligatory

For Peer Review

1
2
3
4 1
5 2
6 2
7
8 3
9
10
11 4
12
13 5
14
15 6
16
17
18 7
19
20 8
21
22
23 9
24
25 10
26
27
28 11
29 12
30
31
32 13
33
34 14
35
36
37 15
38
39
40
41 17
42
43
44 18
45
46
47
48
49 20
50
51 21
52
53
54 22
55
56 23
57
58
59 24
60 25

**Environment and Time as Constraints on the Biogeographical
Distribution of Gibbons**

by

R.I.M. Dunbar

Department of Experimental Psychology, University of Oxford, Oxford, UK

Susan M. Cheyne

Borneo Nature Foundation, Palangka Raya, Indonesia
& Oxford Brookes University, Oxford, UK

Daoying Lan

Guangdong Institute of Applied Biological Resources, Guangzhou 510260, China

Amanda Korstjens

Department of Life and Environmental Sciences, Bournemouth University, Poole, UK

Julia Lehmann

Department of Life Science, University of Roehampton, London, SW15 5JD, UK

G. Cowlshaw

Institute of Zoology, Zoological Society of London, Regents Park,
London NW1 4RY, UK

Abstract

We develop a time budget model for the hylobatid family with the aim of assessing the extent to which their contemporary and historical biogeographic distributions might be explained by ecological constraints. The model uses local climate to predict time budgets, and from this the limiting size of social group that animals could manage at a given location. The model predicts maximum group sizes that vary between 3-15 within the taxon's current distribution, indicating that the combination of their dietary and locomotor styles with the kinds of habitats they inhabit radically constrain group size. Beyond the edges of their current distribution, sustainable group size rapidly tends to zero, although if they had been able to bypass some of these areas, they would have found very suitable habitats in southern India and beyond the Wallace Line. While travel time would be a major constraint on group size at larger group sizes, as it is in great apes, the main factor limiting the gibbon's current distribution is the time they need to spend resting that is imposed on them by the environment. The model also indicates that gibbons would not now be able to survive in regions of central and southeastern China where they are known to have occurred within historical times, perhaps because historical climate change following the Little Ice Age of the C18th made these regions uninhabitable for them. Finally, our results indicate that gibbons have the ecological capacity to live in larger groups than they do, making it unlikely that their adoption of monogamy reflects purely ecological constraints.

Key Words: climate, biogeographic distribution, group size, foraging ecology

1
2
3
4 52
5 53
6 54
7 55
8 56
9 57
10 58
11 59
12 60
13
14
15
16 61
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Highlights:

- Time budgets limit the biogeographic distribution of gibbons and siamang
- Time that has to be allocated to resting seems to be the main constraint
- The limits on group size are not so restrictive as to make monogamy obligatory

For Peer Review

63 Introduction

64 Inability to satisfy nutritional and other demands within a defined time period
65 (usually a 24-hr physiological cycle: Peters 1983) limits both where a species can live
66 and, in social species, how big their groups can be (Dunbar, Korstjens & Lehmann
67 2009). In this respect, time is a central problem for all animals, especially for those
68 that, like primates, are not active 24 hours a day. This insight has led to the
69 development of a series of time budget models designed to understand the role of time
70 in limiting the biogeographical distributions of individual taxa (Dunbar et al. 2009).
71 These models are premised on the fact that behaviour (as the outcome of core
72 physiological processes) is the interface between the environment and the animals'
73 ability to survive (with the latter indexed by the size of group they can maintain as an
74 ecological entity) (see also Marshall, Carter, Rowcliffe & Cowlishaw 2013). There is
75 little that most animals can do to alter their physiology in response to changes in
76 climate or vegetation, but they can, and do, adjust their behaviour and group sizes in
77 response to changing environmental conditions.

78 For species like anthropoid primates that (with obvious the exception *Aotus*)
79 are strictly diurnal due to poor night vision, the time available for satisfying their
80 nutritional requirements is further reduced to the ~12 hours of tropical daylight. In
81 addition, climate and the species' dietary adaptations may force animals to rest
82 ('enforced rest' *sensu* Korstjens et al. 2010), thereby further reducing the length of
83 their active day. The climatic component of this relationship reflects the fact that
84 almost all tropical animals are obliged to cease being active and seek shelter during
85 the hottest part of the day when ambient temperatures in open sunlight exceed their
86 thermoneutral zone (Mount 1979; David-Barrett & Dunbar 2016); the dietary aspect is
87 mainly a problem for folivores for whom gut fermentation of foliage is incompatible

1
2
3 88 with any form of activity because the microbial activity on which it depends is
4
5 89 extremely sensitive to the heat generated by physical activity (van Soest 1982).
6
7

8 90 Intensely social species, like most anthropoid primates, face an additional
9
10 91 problem because the functionality of their relationships, as well as the cohesion of
11
12 92 their social groups, depends directly on the time they invest in grooming with each
13
14 93 other (Lehmann, Korstjens & Dunbar 2007a; Pollard & Blumstein 2008; Dunbar &
15
16 94 Shultz 2010; Dunbar & Lehmann 2013; Sutcliffe, Dunbar, Binder & Arrow 2012;
17
18 95 Dunbar 2018a,b). For both humans and nonhuman primates, an individual's
19
20 96 willingness to give coalitionary aid to another depends directly on the amount of time
21
22 97 they spend engaged in affiliative interaction (Seyfarth & Cheney 1984; Dunbar 1980,
23
24 98 2018a,b; Burton-Chellew & Dunbar 2015). At some point, animals will inevitably run
25
26 99 out of time, and this will ultimately limit where they can live. Ecologists tend to see
27
28 100 this as a constraint imposed by nutrient availability, but the real constraint is time.
29
30
31
32

33 101 In effect, animals face a three-way optimization problem in which they have to
34
35 102 offset the costs of living in social groups (i.e. time investment in social grooming) and
36
37 103 the acquisition of nutrients (specified by the ecological determinants of time required
38
39 104 for feeding and travel, including the fixed costs of fuelling fertility and lactation)
40
41 105 against the benefits of group size as their principal form of predator deterrence.
42
43 106 Predation risk will depend on whether the animal is terrestrial or arboreal and on the
44
45 107 hunting styles of the local predators (the 'landscape of fear') (Shultz et al. 2004;
46
47 108 Shultz & Finlayson 2010; Tolon et al. 2009; Laundré et al. 2014; Coleman & Hill
48
49 109 2014; Riginos 2015; Gallagher, Creel, Wilson & Cooke 2017), but within these
50
51 110 constraints the prey species can adjust group size to offset local predation risk.
52
53
54

55
56 111 Focusing on time budgets is not an alternative to the more conventional socio-
57
58 112 ecological approach that typically seeks to identify correlations between, on the one
59
60

1
2
3 113 hand, group size, population density or behaviour and, on the other hand, climate or
4
5 114 forage quality/distribution (Wrangham, Gittleman & Chapman 1993; Chapman &
6
7
8 115 Chapman 2000; Snaith & Chapman 2007; Clutton-Brock & Janson 2012), or
9
10 116 approaches that explore the efficiency of foraging (e.g. Brockleman et al. 2014).
11
12 117 Rather, a focus on time budgets provides a way of looking at the mechanisms that
13
14 118 underpin the correlations between environmental variables and population level
15
16 119 outcomes (group size and dispersion). Time budgets thus provide insights into the
17
18 120 mechanisms involved in these processes, and, in doing so, emphasize a component of
19
20 121 that mechanism (namely, time) that is invariably overlooked in most studies.
21
22

23
24 122 In addition, a time budget approach allows us to incorporate, directly or
25
26 123 indirectly, a range of other factors and causal relationships that bear on animals'
27
28 124 ability to survive in particular locations. This is important because biological
29
30 125 phenomena are naturally systems-based, with most causal relationships being
31
32 126 multivariate and subject to the influence of feedback loops. Failure to incorporate
33
34 127 these more complex relationships may result in the over- or underestimation of the
35
36 128 importance of particular causal relationships. Time budget models allow us to
37
38 129 integrate a range of variables and relationships into a single, coherent model.
39
40

41
42 130 The formal structure of time budget models is that of a causal chain. The
43
44 131 climate at a given location directly or (via its effect on vegetation quality) indirectly
45
46 132 determines the three core elements of the time budget (feeding, moving and minimum
47
48 133 or 'enforced' rest time). These in turn determine the limiting size of group that
49
50 134 animals can maintain at that location because this depends on how much
51
52 135 'uncommitted' time the animals then have available and the parametric relationship
53
54 136 between time invested in relationships and the resulting quality of those relationships
55
56 137 (as reflected in group coherence). The maximum possible group size predicted by the
57
58
59
60

1
2
3 138 time budget ultimately determines whether or not the species can live at that location
4
5 139 (Dunbar et al. 2009). It is important to appreciate that these models are location-
6
7
8 140 specific: they make explicit predictions about the behaviour of the animals at a given
9
10 141 location, subject to the particular climatic and vegetational conditions pertaining at
11
12 142 that location. They are also necessarily genus-specific because they reflect the
13
14 143 particular dietary, physiological and body size adaptations that characterize a genus.
15
16
17 144 In this respect, a genus is an ecological species.

18
19 145 Climate variables affect both energy demand (energy animals need to maintain
20
21 146 thermoregulation: Mount 1979) and the rate of nutrient intake through the effect
22
23 147 climate has on the quality and digestibility of forage (and hence the amount of forage
24
25 148 that has to be eaten to extract the required amount of nutrients: van Soest 1982). They
26
27 149 also influence both travel time (through their effect on foraging patch size and inter-
28
29 150 patch distances: Janson & van Schaik 1988; Chapman & Chapman 2000) and rest
30
31 151 time (when animals are forced to seek shelter because ambient temperatures exceed
32
33 152 their thermoneutral zone: Mount 1979; David-Barrett & Dunbar 2016). In addition,
34
35 153 the feedback loops that are invariably an important component of biological systems
36
37 154 can arise when group size, in particular, influences travel time (because larger groups
38
39 155 have to travel further if feeding patches are depleted easily – typically, less of a
40
41 156 problem for folivores), and this in turn requires more time to be devoted to foraging to
42
43 157 replace the energy consumption of the additional travel.
44
45
46
47
48

49 158 Time budget models consist of a set of simultaneous equations (one for each
50
51 159 of the core time budget variables) and thus have the form of a linear programming (or
52
53 160 linear optimization) model (Dunbar 2002). The intersection of these equations defines
54
55 161 the limit on group size that a species can maintain at a given location. If any of these
56
57 162 relationships involve nonlinear components, solving the simultaneous equation set to
58
59
60

1
2
3 163 find the maximum group size usually requires numerical rather than analytical
4
5 164 methods (Dunbar 1992a; Dunbar et al. 2009). However, if all relationships are linear,
6
7
8 165 analytical methods based on substitution can be used. The set of climatic parameters
9
10 166 (the primary drivers, with each as a separate X, Z,... axis) and group size (the final
11
12 167 output, or Y-axis, variable) define a multidimensional state space. The isosurface
13
14 168 defined by points where maximum group size drops below the minimum required for
15
16 169 demographic viability and/or predator defence defines the taxon's biogeographic
17
18 170 distribution. So long as the predicted maximum group size lies above these minima,
19
20 171 the species can survive at those locations, with the magnitude of the difference
21
22 172 between the maximum and these minima providing a measure of how much
23
24 173 ecological stress the animals are under.

25
26
27
28 174 These mechanism-based models are at least as successful as conventional
29
30 175 climate (or niche) envelope models at predicting the geographical distributions of the
31
32 176 taxa concerned (Willems & Hill 2009; Korstjens et al. 2010). In addition, they provide
33
34 177 insights into which aspects of the animals' ecology are responsible for limiting their
35
36 178 distribution (Dunbar et al. 2009). This is because time budget models allow us (a) to
37
38 179 determine how much ecological and demographic stress a particular population is
39
40 180 under (Dunbar 1992a; Lehmann et al. 2007a) and (b) to specify why the species is
41
42 181 unable to occupy particular habitats in ways that directly reflect their physiological
43
44 182 adaptations (Dunbar et al. 2009). Neither of these is possible with conventional
45
46 183 climate envelope models, which are essentially simple correlational relationships. In
47
48 184 addition, because time budget models are driven directly by climate, the models can
49
50 185 be used to explore the consequences of past and future climate change for a taxon's
51
52 186 biogeographic distribution in a more nuanced way (Lehmann et al. 2010; Kortsjens,
53
54 187 Lehmann & Dunbar 2010; Bettridge & Dunbar 2012).

1
2
3 188 Time budget models have been published for a number of terrestrial (gelada:
4
5 189 Dunbar 1992a; baboons: Dunbar 1992b; Bettridge, Lehmann & Dunbar 2010; African
6
7 190 great apes: Lehmann, Korstjens & Dunbar 2007b, 2008a) and arboreal (spider
8
9 191 monkeys: Korstjens, Verhoeckx & Dunbar 2006; colobins: Korstjens & Dunbar 2007;
10
11 192 vervets: Willems & Hill 2009; forest guenons: Korstjens, Lehmann & Dunbar 2018;
12
13 193 orangutans: Carne, Semple & Lehmann 2012) primates, as well as one ungulate (feral
14
15 194 goats: Dunbar & Shi 2013).

16
17
18
19 195 Although models have been developed for all three great apes (Lehmann et al.
20
21 196 2007b, 2008a,b; Carne et al. 2012), the small apes (the gibbon family, Hylobatidae)
22
23 197 have yet to be considered. The small apes offer a particularly interesting challenge for
24
25 198 several reasons. First, for an ape, they occupy an unusually exclusive arboreal
26
27 199 terminal branch niche otherwise occupied only by some of the smaller monkeys.
28
29 200 Second, they typically live in very small groups (monogamous pairs plus dependent
30
31 201 offspring) and this raises questions as to why they have opted for this form of social
32
33 202 system. One classic explanation for monogamy in mammals is that females are forced
34
35 203 by their ecology to forage on their own in territories that are too large for a male to be
36
37 204 able to successfully defend more than one female (Wittenberg & Tilson 1980; van
38
39 205 Schaik & Dunbar 1990; Komers & Brotherton 1997; Lukas & Clutton-Brock 2013);
40
41 206 an alternative suggestion is that male parental support allows females to reduce their
42
43 207 energy demand, especially in strongly seasonal habitats or where locomotor costs
44
45 208 prohibit fat storage (Dunbar 1995a,b; Heldstab, van Schaik & Isler 2017). Third, they
46
47 209 have a relatively limited distribution in southeast Asia, and this limited range begs
48
49 210 explanation.

50
51
52 211 Historically, hylobatids probably evolved as a distinct family in southwest
53
54 212 China in the late Miocene, and gradually extended their range down through the
55
56
57
58
59
60

1
2
3 213 Malay peninsula into the islands of the Sunda shelf as opportunities provided by
4
5 214 changes in climate and sea level allowed (Jablonski & Chaplin 2009). Turvey, Crees
6
7
8 215 & Di Fonzo (2015) found that, as late as the 1800s, gibbons were still being recorded
9
10 216 in historical documents as widely distributed throughout southern and central China as
11
12 217 far north as latitude $\sim 35^{\circ}\text{N}$ (Shanxi, Shaanxi and Shangdong provinces). What
13
14 218 explains their disappearance from all but the southwest corner of China (Yunnan and
15
16
17 219 Hainan) remains to be determined, though anthropogenic factors have been suspected
18
19 220 (Fan 2017).

21
22 221 In this paper, we have five main aims. The first is to develop a model of
23
24 222 hylobatid time budgets in terms of climatic variables. Second, we test the model's
25
26 223 validity by using it to predict the taxon's presence and absence in various locations
27
28 224 within continental Asia and the islands of the Sunda Shelf and New Guinea. Third, we
29
30 225 use the model to determine which time budget variables are most responsible for
31
32 226 limiting the taxon's biogeographic distribution. Fourth, we use the model to determine
33
34 227 the possible causes of the historical extinction of gibbon populations in China.
35
36 228 Finally, we use the results to assess how plausible it is that female spacing best
37
38 229 explains hylobatid monogamy.
39
40
41
42
43

231 **Methods**

232 *Gibbon data*

44
45
46
47 233 A database was compiled from the gibbon literature to provide quantitative
48
49 234 data on key demographic, environmental and ecological variables. We identified 77
50
51 235 studies that provide data on at least some of these (see online *Dataset S1*). Of these,
52
53 236 59 provide data on mean group size, 41 provide data on group density or biomass, and
54
55 237 29 provide data on activity budgets and other behavioural or demographic variables.
56
57
58
59
60

1
2
3 238 In respect of activity budgets, we identify five mutually exclusive states (feeding,
4
5 239 moving or travel, resting, social interaction – mainly social grooming – and singing)
6
7
8 240 which, in this sample, between them account for ~100% of time.
9

10 241 Inevitably, a number of issues arise when comparing data from different
11
12 242 studies. One is that studies vary in the definitions they use for individual behaviour
13
14 243 categories; a second is that methods of collecting time budget data vary, both in
15
16 244 intensity (e.g. some researchers use group scans, others focal individual sampling) and
17
18 245 in coverage (the hours of daylight may not be sampled evenly). While these are
19
20 246 certainly issues, past experience developing time budget models suggests that the
21
22 247 impact of these methodological issues is at best modest (see Dunbar & Shi 2013).
23
24 248 More importantly, the main consequence of variations in definition and procedure is
25
26 249 that they increase the error variance in parameter estimates; increased error variance
27
28 250 will simply make it harder to obtain significant results, and will therefore bias the
29
30 251 statistical analysis in favour of the null hypothesis (i.e. no relationship). Ultimately, of
31
32 252 course, the real test is whether our models predict what we see on the ground *despite*
33
34 253 these methodological flaws. We should be less concerned with standardisation of
35
36 254 definitions or methods (though these should always be encouraged) than with whether
37
38 255 models based on them correctly predict what we know to be the case.
39
40
41
42
43

44 256 Where these were given by the original field sources, we also extracted data
45
46 257 on a number of ecological and behavioural variables: the percentage of leaf in the diet
47
48 258 (%Leaf), percentage of fruit in the diet (%Fruit), the density of gibbon groups
49
50 259 (groups/ha), gibbon population biomass (kg/km²), mean day journey length (km),
51
52 260 mean territory size (ha), density of fig trees (Figdens, indexed as trees/ha) and number
53
54 261 of sympatric primate species. Mean day journey length and time spent moving
55
56 262 (transformed into hours spent moving per day) were used to calculate travel speed
57
58
59
60

1
2
3 263 (m/hr). Together, these comprise the set of ecological and covariate variables we will
4
5 264 use in model-building.
6

7
8 265 Data for all sites are provided in online *Dataset S1*.
9

10 266

11
12 267 *Climatic and geophysical data*
13

14 268 Rainfall, altitude and latitude for each study site are those provided by the
15
16 269 cited publications, where these are given. Since latitudinal effects should be
17
18 270 symmetrical about the equator, we transform all latitudes into absolute latitude. Since
19
20 271 individual studies often do not provide all the indices we need, we have sourced all
21
22 272 temperature variables (mean annual temperature, TEMP; mean minimum temperature,
23
24 273 TEMP_{min}; mean maximum temperature, TEMP_{max}; mean daily temperature variation,
25
26 274 TEMP_{var} [difference between mean minimum and mean maximum temperature]; and
27
28 275 the standard deviation of mean monthly temperature, T_{moSD} [an index of seasonality])
29
30 276 from either http://www.globalspecies.org/weather_stations/ (which provides climate
31
32 277 data for individual wildlife reserves) or <http://en.climate-data.org> (which provides
33
34 278 climate data for civic weather stations). Wherever possible, we gave preference to the
35
36 279 first. These climatic variables and the two geographical variables (i.e. latitude and
37
38 280 altitude, both of which are determinants of climate) constitute the set of climatic
39
40 281 variables that we will use as the main independent variables in model-building.
41
42

43
44 282 Previous analyses of weather station data for sub-Saharan Africa have
45
46 283 demonstrated that only three climatic variables are needed to predict evapo-
47
48 284 transpiration (the principal predictor of tropical primary productivity: Rosenzweig
49
50 285 1968; Le Houérou & Hoste 1977; Lo Seen Chong, Mougín & Gastellu-Etchegorry
51
52 286 1993). These are mean ambient temperature, total annual rainfall and an index of
53
54 287 seasonality (Williamson & Dunbar 1998; see also Hill 1999). As in our previous
55
56
57
58
59
60

1
2
3 288 models for African primates, the standard deviation of mean monthly temperature
4
5 289 (TmoSD) is our main index of seasonality. This also allows us to include any effects
6
7 290 due to the fact that, in some contexts, certain times of the year when resource
8
9 291 availability is poor or climate especially challenging may impose limits on what
10
11 292 animals can do (see, for example, Dunbar & Shi 2013).

12
13
14 293 Collinearity between the climate variables in the field site dataset is, in
15
16 294 general, low and within conventionally acceptable levels, the only exception being
17
18 295 that between rainfall and temperature (mainly due to the effect of the cold, dry
19
20 296 habitats in southwest China disproportionately influencing what is otherwise a much
21
22 297 less clearcut relationship). At best, this relationship explains only 20% of the variance,
23
24 298 and as little as 10% if the Chinese habitats are excluded.
25
26
27
28
29

30 *Model-building procedure*

31
32
33 301 For the model-building analysis, we excluded time budget data for one study
34
35 302 on a heavily logged habitat (Johns 1986) because it had an implausibly low estimate
36
37 303 for time spent feeding (8%, >3 standard deviations below the mean for gibbons, and
38
39 304 almost a full SD below the next lowest value) and an implausibly high value for time
40
41 305 spent resting (62%, 2.5 SDs above the overall mean, and a full SD above the next
42
43 306 highest value), two studies (West Garo Hills, NE India: Alfred & Sati 1986, 1990;
44
45 307 Ujung Kulon, Indonesia: P. Kappeler 1984a,b) that had very high values for time
46
47 308 spent feeding (>60%, 3.1 and 3.9 SDs above the mean for gibbons, and >1 SD above
48
49 309 the next nearest value) and unusually low values for resting time, and one (Tanjong
50
51 310 Triang: Ellefson 1974) that had an unusually high value for moving (47%) and low
52
53 311 value for rest (7%). Such high values for feeding and moving and low values for
54
55 312 resting are indicative of gibbons that were not fully habituated.
56
57
58
59
60

1
2
3 313 In developing any model, we face a choice between detail and generality. It is
4
5 314 always possible to build a model that takes every conceivable environmental variable
6
7 315 into account, and hence is 100% accurate in its fit to the data. But in doing so, we
8
9 316 inevitably lose generality: in order to make predictions about where the taxon can
10
11 317 live, we will need to know much more about the particular environmental parameters
12
13 318 of an individual location (e.g. soil type and acidity, inclination and tree composition,
14
15 319 as well as all the larger scale variables like rainfall and temperature). Using general
16
17 320 climate as the basis for a model allows us, at the expense of some loss of accuracy, to
18
19 321 be more general, since relevant climate values can be mapped geographically on a
20
21 322 continental scale from climate models, as well as being projected backwards and
22
23 323 forwards in time.

24
25
26
27
28 324 For the purposes of building a time budget model, we need to determine
29
30 325 taxon-specific equations for just two key variables (feeding and moving time). The
31
32 326 other two main components (enforced resting time and social time) use general
33
34 327 primate equations given, respectively, by Korstjens et al. (2010) and Lehmann et al.
35
36 328 (2007a). For the purposes of the present model, we shall take time devoted to singing
37
38 329 as a constant (at the mean observed value of 4.96%).

39
40
41
42 330 Enforced resting time, as defined by Korstjens et al. (2010), specifies the
43
44 331 minimum amount of time that has to be spent resting as a consequence of high
45
46 332 ambient temperatures (to avoid thermal overload or excessive heat loss) and for the
47
48 333 digestion of leaves (as a function of climatic conditions and the species' dietary
49
50 334 physiology). Enforced resting time differs from observed resting time (time spent
51
52 335 inactive, as recorded by field observers) in that observed rest time consists of enforced
53
54 336 resting time plus uncommitted time ("free rest") (Korstjens et al. 2010).

55
56
57
58 337 The time during the day that animals have left over after making their habitat-

1
2
3 338 specific allocations to feeding, moving, enforced rest and singing (subject to any
4
5 339 feedback loops in the equation set) gives us the amount of time in the day that could
6
7
8 340 be devoted to social interaction (grooming). Because grooming time correlates with
9
10 341 group size (Dunbar 1992; Lehman et al. 2007; Dunbar & Lehmann 2014), we can use
11
12 342 this relationship to specify the maximum size of group that the animals could maintain
13
14
15 343 as a coherent social entity at that location. This does not mean the animals *have* to
16
17 344 devote all this time to social interaction: it simply sets the upper limit, and hence the
18
19 345 upper limit on social group size.

21 346 The social time equation reflects the fact that social grooming is the principal
22
23 347 bonding mechanism for primate social groups and increases linearly with group size
24
25
26 348 across primates as a whole (Dunbar 1991; Lehmann et al. 2007; Dunbar & Lehmann
27
28 349 2014). As such, it represents the time investment in social interaction that is necessary
29
30 350 to maintain a group's cohesion through time so as to prevent it disintegrating (Dunbar
31
32 351 1992a, 2012). It is worth noting that this increase in time devoted to grooming with
33
34 352 group size does not mean that animals in large groups groom with more individuals.
35
36
37 353 Typically they do not. Rather, as group size increases, animals invest increasingly in
38
39 354 the handful of core grooming partners that act as their primary coalition partners
40
41 355 within the group because these act as their primary defence against the stresses of
42
43 356 living in a group (Kudo & Dunbar 2001; Dunbar 2012, 2018a).

46 357 As in previous models, we first examined all relevant pairwise plots visually to
47
48 358 determine whether any factors correlate nonlinearly with the main time budget
49
50 359 variables. There were no nonlinear relationships. We then ran backward stepwise
51
52 360 linear regression models with the full set of climatic and ecological variables as
53
54 361 predictors of each of the two dependent variables (feeding and moving time). The
55
56 362 criterion used in selecting the final equation in each case was based on maximising
57
58
59
60

1
2
3 363 the proportion of variance explained, subject to the requirement that the overall
4
5 364 equation and all its individual predictor variables were significant.
6
7

8 365 Because siamang are more than twice as large as other gibbons, siamang
9
10 366 feeding time was first corrected to a gibbon-equivalent (i.e. what it would need to do
11
12 367 if it was a conventional gibbon) by rescaling it by the ratio of the metabolic body
13
14 368 weights of the two taxa (i.e. $5.5^{0.75}/10.75^{0.75}=0.605$). This allows us to use all the data
15
16 369 without having to include an additional factor for taxon or body mass. None of the
17
18 370 other gibbon species differed significantly in body weight (see *Dataset S1*).
19
20

21 371 In extracting these equations from the data, we have ignored the possible
22
23 372 influence of phylogeny. As in previous analyses (see Dunbar et al. 2009), we do so
24
25 373 principally because almost all of the variables we are concerned with are behavioural
26
27 374 rather than anatomical, and are likely to be influenced more heavily by local
28
29 375 environmental conditions than by biological inheritance, and thus typically have
30
31 376 phylogenetic signals that are close to zero (see Kamilar & Cooper 2013). In effect, we
32
33 377 treat all the hylobatids as belonging to a single ecological species. Inter-population
34
35 378 analyses of just this kind have previously been successfully used to explore aspects of
36
37 379 the behaviour of gibbons (song function: Cowlshaw 1992, 1996). It is perhaps
38
39 380 important to note that we are not seeking to describe the “typical” behaviour of
40
41 381 individual gibbon species: rather, our concern is to understand the overall pattern of
42
43 382 behavioural flexibility and the effects of climate on biogeography across the whole
44
45 383 taxon. For convenience, we adopt the convention of referring to the entire group as
46
47 384 hylobatids, and differentiate between siamang and all other gibbon species only on the
48
49 385 basis of body size.
50
51

52 386 We then use this set of equations to determine maximum group size for any
53
54 387 given location. Since all the equations in the present case are linear, we use analytical
55
56
57
58
59
60

1
2
3 388 methods and solve directly by substitution:
4

5 389
$$N_{Max} = (100 - (\text{Feed} + \text{Move} + \text{Rest}_{\text{Enf}})) / (\text{grooming equation})$$

6
7

8 390 This value is the limiting group size (the maximum ecologically tolerable group size,
9
10 391 N_{max}).

11
12 392 In order to exhibit the biogeographical implications of the model in graphable
13
14 393 form, we present the main results as 3-dimensional surface plots of maximum group
15
16 394 size plotted against (a) absolute latitude and altitude and (b) mean annual rainfall and
17
18 395 mean annual temperature. Because a relatively large number of environmental and
19
20 396 behavioural variables are involved in the basic model, we need to convert all other
21
22 397 variables into functions of just the two variables used for each graph. We do this by
23
24 398 using backwards stepwise regression modelling to derive equations for these variables
25
26 399 from the data given in online *Dataset S1*. The resulting equations are given in Table 2.
27
28 400 Of these, latitude, altitude and temperature are well known to be systematically
29
30 401 related.
31
32

33
34
35 402 All statistical analyses and graphs were executed in SPSS v.23.
36
37
38 403

39 404 *Testing the model*

40
41 405 We test the model by evaluating its ability to predict the presence versus
42
43 406 absence of gibbons at a range of locations across the Indian subcontinent, the Indo-
44
45 407 China peninsula, Indonesia and the Malay Archipelago, New Guinea and mainland
46
47 408 China. For these purposes, we selected a number of locations of known altitude and
48
49 409 latitude, which had climatic data available in www.en.climatedata.org. The locations
50
51 410 and their climatic and geographical data are given in online *Dataset S2*. Since these
52
53 411 analyses predict where hylobatids can and cannot survive, they also allow us to make
54
55 412 inferences about the biogeographic distribution of the taxon. In most cases, we
56
57
58
59
60

1
2
3 413 selected major cities for these purposes, since our question is whether or not gibbons
4
5 414 could live in the general area, not whether they live at a particular location. These data
6
7 415 also allow us to assess whether gibbons could now live in those provinces of China
8
9 416 where they were recorded historically by Turvey et al. (2015), and hence determine
10
11 417 whether anthropogenic factors or climate change might have been responsible for
12
13 418 their extinction. For these purposes we selected one site at random roughly in the
14
15 419 center of each of the provinces of modern China.
16
17
18
19
20

420

421

Results

The model

21
22
23
24 422
25
26 423 Table 1 lists the best fit equations for feeding and moving that were generated
27
28 424 by the data, as well as the generic primate equations for enforced resting time and
29
30 425 social time. The causal relationships involved are summarized in Fig. 1. For reference,
31
32 426 alternative “next best” significant equations for feeding and moving time selected by
33
34 427 the backwards stepwise regression are given in the *Online Supplementary Material*
35
36 428 (Table S1). Both feeding and moving time increase as habitats become climatically
37
38 429 more stressful (low rainfall, low temperature, high altitude, high latitude). Although
39
40 430 group size has no effect on feeding time, it has a weak positive effect on moving time
41
42 431 even despite the extremely limited variation in hylobatid group size (Fig. 2). Since
43
44 432 moving time is the main limiting factor for great apes, and severely limits their
45
46 433 biogeographic distribution and group size (Lehmann et al. 2007b, 2008a,b), we ran a
47
48 434 separate regression model with just latitude and group size as predictors (Eqn. 2a). It
49
50 435 is of significance that, despite an order of magnitude difference in the range of group
51
52 436 sizes, the slope coefficient for group size in gibbons is very similar to that for the
53
54 437 chimpanzee time budget model (3.08 vs 2.59, respectively).
55
56
57
58
59
60

1
2
3 438 Across the sample of study populations, percentage of fruit in the diet is
4
5 439 determined mainly by ambient temperature (Table 2). In contrast, percentage of leaf
6
7 440 in diet increases with the level of environmental stress as reflected in rainfall, altitude
8
9 441 and temperature variance (Table 2). Although day journey length was not predicted by
10
11 442 any of the climatic variables or by fig tree density, territory size is predicted with a
12
13 443 very high r^2 by a complex of climatic variables (Table 2), suggesting that territory
14
15 444 sizes get larger as environmental stress increases.
16
17
18

19 445 The next step is to use the time budget equations to predict maximum
20
21 446 ecologically possible (i.e. limiting) group size, N_{max} , for the sampled populations. To
22
23 447 do this, we interpolate the time available for social interaction (i.e. that remaining after
24
25 448 removing the time predicted for feeding, moving and enforced rest at the site, plus the
26
27 449 constant for singing) into Eqn. (5). For this analysis, siamang predicted feeding time
28
29 450 is rescaled back to siamang body mass by reversing the transformation used to convert
30
31 451 their feeding time to gibbon-equivalents.
32
33
34

35 452 Fig. 3(a) plots maximum tolerable group size, N_{max} , against observed mean
36
37 453 population group size for all the gibbon and siamang populations in our sample. The
38
39 454 horizontal line demarcates a group size of 2.96 individuals, representing the minimum
40
41 455 group size for demographic viability (i.e. demographic stability defined by a
42
43 456 population growth rate of $r=0$: this requires two surviving offspring over an average
44
45 457 gibbon 17-year reproductive lifespan, and would equate to a group with two adults
46
47 458 plus $4/17 = 0.12$ births per year, hence an average of ~ 0.96 dependent offspring aged
48
49 459 < 8 years at any given time, taking 8 to be the age at dispersal and thus the range of
50
51 460 individuals who would be observed in a group). (This value assumes minimal
52
53 461 mortality in immature gibbons, which seems broadly to be the case; assuming 50%
54
55 462 mortality up to puberty, a figure typical of terrestrial baboons, would only increase
56
57
58
59
60

1
2
3 463 minimum viable group size to 2 adults + 1.5*0.96 living offspring = 3.44.) The
4
5 464 dashed diagonal line marks the line of equivalence (N_{max} is equal to observed mean
6
7
8 465 population size).

9
10 466 Three points may be noted. First, time budget models specify that group sizes
11
12 467 should not exceed the predicted maximum size (hence all data points in Fig. 3a should
13
14 468 lie above the dashed line). In the present case, 80.5% of the 41 populations have a
15
16 469 predicted maximum group size that is larger than their observed mean group size.
17
18 470 This is broadly encouraging as far as model fit is concerned, but suggests that
19
20 471 something else is needed to account for the exceptions. Second, no population has
21
22 472 $N_{max} > 15$ individuals. This suggests that hylobatids are under rather greater ecological
23
24 473 constraint than many other Old and New World monkeys (and African great apes),
25
26 474 where limiting group sizes are typically > 20 . Third, notwithstanding this, it is clear
27
28 475 that many (but not all) populations could, on ecological grounds, live in much larger
29
30 476 groups than they actually do. Such groups would inevitably be multi-female.

31
32
33
34
35 477 One likely reason why some populations might have maximum predicted
36
37 478 group sizes below their observed group sizes is the length of the active day. Gibbons
38
39 479 are well known to retire early, often several hours before dusk (Raemaekers 1979;
40
41 480 Chivers 1984; Palombit 1997; Fei, Zhang, Yuan, Zhang & Fan 2017), a behaviour
42
43 481 that is quite unusual for most primates. As a result, the sampled activity budgets are
44
45 482 based on an active day that averages 9.3 ± 0.89 hrs (range 8.1-10.6, $N=9$ sites), despite
46
47 483 the fact that daylength in tropical habitats is ~ 12 hrs. Progressively extending the
48
49 484 length of the active day in units of 5% suggests that an active day that is 15% (84
50
51 485 min) longer than the observed mean would be enough to lift all but the three lowest
52
53 486 siamang populations above the demographic viability threshold (Fig. 3b). (This would
54
55 487 increase maximum N_{max} for the other populations only to ~ 20 .) Unfortunately, data on
56
57
58
59
60

1
2
3 488 the length of the active day are not available for any of the populations with
4
5 489 $N_{max} < 2.96$. However, an extra 84 min would only increase the mean value of day
6
7 490 length to 10.7 hrs, virtually identical to the longest observed day length in our dataset
8
9 491 (10.6 hrs in a siamang population: *Dataset S1*).

12 492 We ran a sensitivity analysis of the model using the alternative equations for
13 493 feeding and moving time from Table S1. To do this, we altered one of the equations at
14 494 a time in the original model from Table 1, keeping all the other equations as in the
15 495 original model. We also ran a model combining the alternative moving time equation
16 496 with the two most extreme alternative feeding time equations. Predicted maximum
17 497 group size for the individual populations in the study site database determined using
18 498 these alternative equations are highly correlated with those determined using the
19 499 original model (Table S2; Fig. S1). The alternative moving time equation increases
20 500 the largest maximum group size from ~15 to ~25, but does not substantively change
21 501 any of the results. This suggests that the Table 1 model is quite robust. Similarly,
22 502 decreasing the slope coefficient for group size in the moving time equation to 2.59
23 503 (the value for the chimpanzee model) increases the largest N_{max} to ~18, while
24 504 increasing it to 5.0 decreases N_{max} to ~9, but does not alter the broad pattern or the fact
25 505 that most mean population group sizes are comfortably below N_{max} .

26 506 To provide some indication as to how environmental parameters influence
27 507 hylobatid biogeography, Fig. 4 plots predicted maximum group size as a function of
28 508 (a) latitude and altitude and (b) rainfall and temperature. To produce these graphs, we
29 509 have used the equations given in Table 2 to reduce all the climatic variables in the
30 510 model to the two indices of interest in each case. This can be expected to increase
31 511 error variance, with a consequential tendency for reversion to the mean, but the results
32 512 provide us with an indication of how maximum group size is likely to vary across

1
2
3 513 habitats. The results suggest that hylobatids do best (i.e. are able to sustain larger
4
5 514 groups) in cooler, high altitude, high rainfall habitats at low latitudes (i.e. near the
6
7 515 equator). When rainfall is less than ~1500mm per annum, or at latitudes above ~20°
8
9
10 516 (in effect, outside the Tropics), hylobatids are unable to maintain minimally viable
11
12 517 groups ($N \approx 3$ individuals) unless they are living at altitudes above ~1000m (i.e. under
13
14 518 cooler conditions) (see also Turvey et al. 2015). This last prediction is confirmed by
15
16
17 519 the fact that the northern Chinese populations are at significantly higher altitudes than
18
19 520 all the other hylobatid populations (mean altitudes: 1750.0 ± 783.8 SD m vs
20
21 521 401.2 ± 409.5 SD m asl, $N=14$ and 50 respectively; $F_{1,62}=76.2$, $p < 0.0001$).

22
23
24 522

25
26 523 *Testing the model*

27
28 524 To explore the model's ability to predict hylobatid biogeography, we used a
29
30 525 sample of locations within the current hylobatid biogeographic range in southeast
31
32 526 Asia and a sample of locations on the Indian subcontinent, mainland China and the
33
34 527 islands of the Malay archipelago outside the current hylobatid range. In addition, we
35
36 528 have included one representative site in each of the Chinese provinces where gibbons
37
38 529 were recorded as occurring prior to 1800 AD (see Turvey et al. 2015). Note that, for
39
40 530 the latter cases, the climate data are current, not historical, values. In each case, we
41
42 531 predict maximum group size using current local climate
43
44
45

46
47 532 Fig. 5 plots the mean and range of maximum group size predicted by the
48
49 533 model for each of these sites grouped by geographical location. Predicted maximum
50
51 534 group sizes average 6.6 ± 4.0 SD for 18 sites within the current hylobatid range. In
52
53 535 contrast, sites in northeast India and Bangladesh to the west of the Brahmaputra
54
55 536 (which forms the northwest boundary of the current gibbon range) average 0.3 ± 0.5 SD
56
57
58 537 ($N=7$ sites), as is the case for locations further west in northwest India (mean
59
60

1
2
3 538 1.6±2.2SD, N=7 sites) (see also Fig. S2). This suggests that, even in the absence of
4
5
6 539 the physical barrier of the Brahmaputra river system, the taxon's ability to expand its
7
8 540 range westwards has been limited by substantial tracts of land it would have been
9
10 541 unable to cross. Had gibbons been able to bypass these two barriers, they would have
11
12 542 found very congenial habitats in the southern parts of the subcontinent (e.g. the Nilgiri
13
14 543 Hills and adjacent ranges in the Deccan, and in Sri Lanka), where they would have
15
16 544 been able to support groups as large as 8.7±5.1SD (N=7 sites). The island habitats
17
18 545 across the Wallace Line (specifically, the Celebes and Papua New Guinea) provide
19
20 546 equally gibbon-friendly habitats (mean group size 10.3±1.1SD, N=2 sites), but it
21
22 547 seems that, as with most other primates (Brandon-Jones 1998), the gibbons were
23
24 548 never able to traverse this major sea barrier. Increasing the length of the available day
25
26 549 by 15% (as in Fig. 3b) does not change the overall pattern, or make regions such as
27
28 550 northern India or central China any more habitable for gibbons.

32
33 551 More puzzling, however, is the fact that central and east Java (including
34
35 552 nearby islands like Bali) provide very suitable habitats (mean N_{max} 10.4±1.2SD, N=7
36
37 553 sites) yet have no gibbon populations. Climatically at least, these habitats seem to be
38
39 554 at least as suitable as those in west Java where gibbons do occur (mean N_{max}
40
41 555 10.2±1.9SD, N=7 sites). In contrast, gibbons would now find it impossible to survive
42
43 556 in either the central and southeastern provinces of China where they were recorded as
44
45 557 living prior to 1800 AD (mean 0.0±0.0SD, N=14 sites) or, perhaps less surprisingly,
46
47 558 the provinces of northern China where they did not occur historically (mean
48
49 559 1.3±3.6SD, N=7). This contrasts with locations in southwest China (Yunnan province
50
51 560 and Hainan Island) that currently do support gibbon populations, for which the model
52
53 561 predicts viable group sizes (mean N_{max} = 4.3±1.7SD, N=3 sites).
54
55
56
57
58
59
60

1
2
3 563 *What limits hylobatid distribution?*
4

5 564 An important feature of time budget models is that they allow us to determine
6
7
8 565 which aspect of the animals' biology is the principal constraint on their ability to
9
10 566 occupy habitats, and hence what actually limits their biogeographic distribution. In
11
12 567 turn, this tells us something about the taxon's risk of extinction under climate change.
13
14 568 To explore this, we plotted the predicted time required for feeding, moving and
15
16 569 enforced rest for the *Dataset S2* locations (Fig. 6). Neither feeding nor moving differ
17
18 570 consistently between regions where hylobatids are present vs not present, suggesting
19
20 571 that neither of these is a major constraint (except, in the limit, the effect of group size
21
22 572 on moving time at much larger group sizes); in contrast, resting is significantly higher
23
24 573 in regions where they are absent and seems to account for the fact that they do not
25
26 574 have sufficient time for social interaction to maintain groups of any significant size.
27
28
29

30 575 It thus seems that the main problem lies in the extent to which climatic
31
32 576 variables oblige them to rest. This is confirmed by an analysis of observed versus
33
34 577 predicted time budget allocations in the sample study sites. Fig. 7a plots the ratio of
35
36 578 observed feeding and resting time divided by the values predicted by the model
37
38 579 equations in Table 1 for those populations in the field study dataset (*Dataset S1*) for
39
40 580 whom predicted maximum ecologically tolerable group size is either less than or
41
42 581 greater than 5 (the upper limit for observed mean population group size). For these
43
44 582 purposes, predicted rest is enforced rest time, while observed rest is total rest time.
45
46 583 Feeding time is close to that predicted in both cases, as should be the case. In contrast,
47
48 584 observed rest time is significantly less than the minimum predicted ('enforced' rest) in
49
50 585 populations where N_{max} is less than 5, whereas populations where N_{max} exceeds 5
51
52 586 typically have surplus rest time capacity. The strength of this effect is clear from Fig.
53
54 587 7b which plots the difference between observed and predicted rest against the
55
56
57
58
59
60

1
2
3 588 difference between observed group size versus predicted N_{max} for the populations.
4
5 589 The correlation is highly significant ($r=-0.847$, $p<0.0001$).
6
7
8 590

591 **Discussion**

592 The time budget model for hylobatids that we develop here is broadly similar
593 in form to the models developed for 10 other primate genera. It works well for
594 contemporary populations, predicting presence in most cases where gibbons occur and
595 absence where they do not (Fig. 5). In other cases, it is clear that physical barriers
596 (e.g. habitat conditions in northeast India or the Wallace Line to the southeast) have
597 prevented gibbons expanding into habitats where they could do well. It is important to
598 remember that time budget models predict the largest groups that a taxon can maintain
599 at a given site (in effect, the carrying capacity), not necessarily the actual size of
600 group. They simply set the upper limit defined by local environmental conditions.
601 Since, for primates in general, fertility is almost always adversely affected by
602 increasing group size (van Schaik 1982; Dunbar 2018a; Dunbar, MacCarron & Shultz
603 2018a), animals will generally try to minimize group size in any given location to the
604 extent that this is compatible with the constraints imposed by the local predation risk
605 (Dunbar et al. 2009; Dunbar et al. 2018a).

606 The fact that hylobatids commonly live in groups that are smaller than those
607 allowed by the local ecology has two important implications. First, it implies that,
608 while the model certainly tells us that gibbons are under greater ecological constraint
609 than most Old World monkeys and African great apes, the level of ecological stress is
610 not sufficiently high to force them to live in groups as small as those they actually live
611 in. While it is true that some populations have time budgets with little or no spare
612 capacity, many do not (Fig. 7). This conclusion is supported by the fact that, unlike

1
2
3 613 most other monkeys and apes, hylobatids commonly go to their night rest mid-
4
5 614 afternoon (Raemaekers 1979; Srikosamatara 1984; Palombit 1999; Fei et al. 2017),
6
7
8 615 implying that they are not under significant time pressure. Second, it suggests that
9
10 616 predation risk must be low for hylobatids; group size evidently does not provide
11
12 617 gibbons with a significant anti-predator advantage, so they can afford to minimize
13
14 618 group size in order to maximize fertility (see Dunbar et al. 2009; Bettridge, Lehmann
15
16 619 & Dunbar 2010; Dunbar, MacCarron & Robertson 2018b). Indeed, in comparison to
17
18
19 620 almost all other primates, cases of predation are conspicuous by their absence from
20
21 621 the gibbon literature (Reichard 2003).

22
23
24 622 This calls into question the claim that gibbons are monogamous because
25
26 623 females are forced by their ecology to forage on their own rather than in groups (van
27
28 624 Schaik & van Hooff 1983; Rutberg 1985; Komers & Brothertpn 1997; Brotherton &
29
30 625 Komers 2003; Lukas & Clutton-Brock 2013). Contrary to this claim, it seems that
31
32 626 most populations could in fact maintain larger groups (range 5-12 animals), especially
33
34 627 in the more southerly parts of the Malay archipelago. Groups of this size could easily
35
36 628 include 2-4 reproductive females (across all primates, reproductive females form a
37
38 629 very consistent 30-35% of the group: Dunbar et al. 2018a,b). Even if females lived
39
40 630 alone, it would be possible to have more of them within a gibbon group's current
41
42 631 territory, and males would have access to more than one reproductive female if they
43
44 632 pursued a roving male strategy (much as orang utan males do) (see also van Schaik &
45
46 633 Dunbar 1990; Bartlett 2009). This implies, as suggested by van Schaik & Dunbar
47
48 634 (1990) and Opie, Atkinson, Dunbar & Shultz (2014), that monogamy must have
49
50 635 evolved in response to factors other than ecological constraints.

51
52
53
54
55
56 636 One of the main benefits of the time budget model approach is that it provides
57
58 637 insights into the behavioural and physiological constraints that limit a taxon's ability
59
60

1
2
3 638 to successfully occupy different kinds of habitats. As with previous primate time
4
5 639 budget models, feeding time is strongly influenced by predictors of habitat quality: in
6
7 640 the hylobatid case, the main determinants of feeding time (rainfall, altitude and
8
9 641 latitude, with the latter two both being core predictors of temperature) are all variables
10
11 642 that influence both tree species composition and the nutritional quality of vegetation
12
13 643 (see also Marshall & Leighton 2006). As food quality falls, animals will need to spend
14
15 644 longer feeding to meet their nutritional requirements.

16
17
18
19 645 In contrast, time spent moving remains largely independent of the climatic
20
21 646 variables, being affected only by absolute latitude and group size. Models for African
22
23 647 genera have also noted that moving time tends to be independent of environmental
24
25 648 variables and is often close to being a constant (Dunbar et al. 2009), mainly because,
26
27 649 in order to save time for other more pressing activities, animals respond by increasing
28
29 650 travel speed rather than increasing travel time when they need to travel further
30
31 651 (baboons: Dunbar 1992a). There is some suggestion that gibbons also do this: travel
32
33 652 speed increases as climatic conditions deteriorate (i.e. when temperatures are high and
34
35 653 latitude is low, although the effect is weak and not significant (Fig. S3). However,
36
37 654 since there is inevitably a limit on the speed at which animals can travel, at some point
38
39 655 animals simply have to devote more time to moving. The substantive issue for day
40
41 656 journey length (and hence moving time) is, at least for non-folivores (Snaith &
42
43 657 Chapman 2007), group size: the area that has to be covered to allow every animal in
44
45 658 the group to meet its nutritional requirements is inevitably a linear function of the
46
47 659 number of animals in the group, forcing the group to travel further each day (Dunbar
48
49 660 et al. 2009). The great apes are especially susceptible to the effects of group size on
50
51 661 moving time, and it is this that is mainly responsible for chimpanzees' fission-fusion
52
53 662 form of sociality (Lehmann et al. 2007b, 2008a,b). It is significant that, despite the
54
55
56
57
58
59
60

1
2
3 663 very limited variance in the size of their groups, the group size slope parameter for the
4
5 664 gibbons is similar to that for chimpanzees. For groups as small as those found in
6
7
8 665 gibbons, the impact of this effect will, of course, be modest; but it does mean that
9
10 666 when groups are larger than ~10 the impact will rapidly becomes prohibitive (as is the
11
12 667 case for African great apes: Lehmann et al. 2007b [Fig. 7]).

13
14 668 However, it seems to be enforced resting time that is the main constraint for
15
16 669 hylobatids (Figs. 6 and 7). Since this is mainly affected by ambient temperature, it
17
18 670 might explain why social groups get larger in northern populations irrespective of
19
20 671 species (Fig. 8). Ultimately, however, the northern extension of hylobatids is
21
22 672 constrained by the fact that feeding and moving time are also positive functions of
23
24 673 latitude (and hence, in effect, declining temperatures). A comparison of the slope
25
26 674 parameters for latitude in Tables 1 and 2 indicates that the additive effects of the
27
28 675 latitude coefficients for feeding and moving time increase at a combined rate that is
29
30 676 ~4 times the savings in terms of the slope for enforced rest ($0.363 + 0.462 = 0.825$ vs
31
32 677 $1.33 \times 0.158 = 0.210$), so that time budgets become progressively squeezed as animals
33
34 678 occupy increasingly high latitude habitats. At the latitude of Mt Wuliang in Yunnan
35
36 679 Province, latitude has added a net 15 percentage points to the time budget.

37
38 680 There is a widespread perception that *Nomascus* populations live in larger
39
40 681 groups than all other hylobatids, hence the fact that they have polygamous groups. Up
41
42 682 to a point, this is true; however, this is not true for all *Nomascus* populations (Fig. 8).
43
44 683 Moreover, *Hoolock* populations also have large groups at high latitudes without these
45
46 684 being polygamous. Mean population group size is in fact a cubic function of absolute
47
48 685 latitude (Fig. 8). Group size does increase steadily up to around 20°N (roughly the
49
50 686 latitude of Chaing Mai in northern Thailand), but then it levels off only to drop
51
52 687 precipitately after latitude 25°N (roughly the latitude of Mt Wuliang in southwestern
53
54
55
56
57
58
59
60

1
2
3 688 China). This decline at very high latitudes is suggestive of populations living at the
4
5 689 limits of their range, and indeed mean group size at these highest latitude populations
6
7
8 690 is only just above the minimum for demographic viability.
9

10 691 It is evident from Fig. 3(b) that siamang incur an additional cost due to the fact
11
12 692 that their larger body size imposes a higher feeding time demand. A convincing
13
14 693 explanation for their larger body size remains elusive, especially given the fact that
15
16
17 694 their range overlaps that of other gibbons and the two taxa are often sympatric. As a
18
19 695 result, however, they are often under significant time budget pressure. There is some
20
21 696 evidence to suggest that they make time budgeting adjustments that might be
22
23 697 sufficient to reduce this pressure: for the limited sample available, it seems that
24
25
26 698 siamang devote less time each day to both singing and social interaction than gibbons
27
28 699 do, and they have a longer active day (Table 3). For this limited sample, the net gain
29
30 700 for siamang is equivalent to increasing the length of the active day by ~17% (the
31
32 701 combined effect of a longer active day adjusted for less time spent socialising and
33
34 702 singing) while only having to increase the actual length of the active day by ~10%.
35
36
37 703 That is sufficient to just lift the lower siamang populations in Fig. 3(b) into the
38
39 704 minimum viability zone. Nonetheless, siamang populations are clearly under more
40
41 705 ecological pressure than is typical of the other gibbons (Fig. 3).
42
43
44

45 706 Gibbons share with the great apes a dietary physiology specialized for
46
47 707 frugivory; indeed, their feeding time equation is very similar to that of the gorilla,
48
49 708 though their size precludes their being able to survive on low quality folivorous
50
51 709 fallback foods in the way gorillas can. Although the larger-bodied siamang exhibit
52
53 710 some capacity in this direction (Raemaekers 1979; Palombit 1995), their ability to
54
55 711 resort to a heavily folivorous diet is likely to be considerably less than the gorilla's
56
57
58 712 simply because of their smaller body size. Most years aside, fruits invariably have a
59
60

1
2
3 713 more patchy distribution than leaf, and the travel demands imposed by this may be
4
5 714 part of the problem that affects gibbon time budgets, and hence limits group sizes.
6

7
8 715 Gibbon ecology appears to be quite tightly defined: their geographical range is
9
10 716 surprisingly limited. They are currently confined to the Indochina peninsula (bridging
11
12 717 out into southwest China) and the associated islands of the Malay archipelago
13
14 718 (Sumatra, Java and Borneo). They appear to have been prevented from extending
15
16 719 southeastwards by the Wallace Line (despite the fact that the islands beyond would be
17
18 720 perfectly habitable for them: Fig. 5), while their capacity to encroach into the Indian
19
20 721 subcontinent has been limited by the climate west of the Brahmaputra (Fig. 6).
21
22 722 Nonetheless, within their core distribution they seem to be under much less ecological
23
24 723 constraint than we might have anticipated given the female-dispersion explanation for
25
26 724 their adoption of a monogamous mating/social system.
27
28
29

30
31 725 Historically, gibbons are known to have occurred at much higher latitudes in
32
33 726 China than they do now. The model suggests that current climate in these areas makes
34
35 727 these habitats completely unsuitable for gibbons. Although the historical extinction of
36
37 728 these northern populations has been attributed to anthropogenic factors (Fan 2017), in
38
39 729 fact these habitats would not support gibbons now even if there were no humans
40
41 730 living there. There are only three plausible explanations for the disappearance of these
42
43 731 populations. One is that the populations in question had evolved novel adaptations to
44
45 732 these environments, such that the slope parameters on their feeding and/or moving
46
47 733 equations were radically different from those for gibbons from further south, but that
48
49 734 humans caused their extermination nonetheless. However, to suggest that these
50
51 735 northern gibbons were doing something completely different to all other New and Old
52
53 736 World anthropoid primates as well as other gibbons implies that they could not have
54
55 737 been conventional primates, and that should be, at best, an explanation of last resort.
56
57
58
59
60

1
2
3 738 A second possibility is that humans had released animals in these areas in order to
4
5 739 have them as exotics (something that humans seem to be especially prone to do), even
6
7 740 though the habitats were not really suitable for them; as a result, the populations went
8
9 741 extinct once they were no longer being replenished (a situation not too dissimilar to
10
11 742 that of *Macaca sylvanus* on Gibraltar). This is a possibility, but the fact that gibbons
12
13 743 seem to be unable to survive there at all would imply the need for constant
14
15 744 replenishment, and this seems unlikely over such a wide area and such a long time
16
17 745 period. The third option is that the climate has changed in the past 300 years in ways
18
19 746 that would have led to the inevitable extinction of these populations (with or without
20
21 747 the assistance of humans).
22
23
24
25

26 748 A potential culprit in the latter respect is the Little Ice Age of 1645-1715,
27
28 749 which resulted in average world temperatures falling by $\sim 1^{\circ}\text{C}$. In China, this resulted
29
30 750 in a significant increase in climatic seasonality. The cultivation of Mediterranean-type
31
32 751 citrus crops was abandoned in Jiangxi Province (one of the provinces that historically
33
34 752 harboured gibbons: Turvey et al. 2015) after the 1750s, despite their having been
35
36 753 cultivated there for many centuries (Reiter 2000). Guangdong (another of the
37
38 754 provinces where gibbons have been documented) experienced a particularly cold, dry
39
40 755 spell, accompanied by an unusually high frequency of typhoon strikes, after the Little
41
42 756 Ice Age (Liu, Shen & Louie 2001). Ameca y Juárez, Mace, Cowlshaw, Cornforth &
43
44 757 Pettorelli (2013) have shown, for mammals as a whole and primates in particular, that
45
46 758 high frequencies of cyclones and droughts correlate with elevated extinction risk
47
48 759 (indexed by the number of taxa classified as “threatened”), with the southeast corner
49
50 760 of China being especially prone to this effect. Turvey et al. (2015) noted that there
51
52 761 was increasing fragmentation of these populations from 1700 onwards (with a marked
53
54 762 upturn in fragmentation from around 1900), with a 50-100 year lag to last reported
55
56
57
58
59
60

1
2
3 763 occurrence. This is indicative of populations struggling to survive.
4

5 764 A comparison of the distribution of climatic variables in southwest China
6
7
8 765 (where gibbons live now) and central and southeast China (where they lived
9
10 766 historically) suggests that the only climate variables in which these two regions differ
11
12 767 significantly are $TEMP_{moSD}$ (standard deviation of mean monthly temperature across
13
14 768 the year, an index of seasonality) and annual rainfall (Table 4). (We do not consider
15
16 769 latitude or altitude since these cannot have changed historically.) This may well
17
18 770 reflect shifting patterns in the latitudinal distribution of the monsoon circulation,
19
20 771 allowing an extension of a more seasonal monsoon climate further into mainland
21
22 772 China after the Little Ice Age. A 65% increase in rainfall and a dramatic reduction in
23
24 773 seasonality would be required to allow gibbons to survive in the southeastern
25
26 774 provinces as well as they currently do in the southwest.
27
28
29

30 775 The bottom line seems to be that even if anthropogenic factors have been
31
32 776 important in the final demise of these populations (as both Turvey et al. 2015 and Fan
33
34 777 2017 have suggested), these populations' sensitivity to anthropogenic factors is likely
35
36 778 to have been exacerbated by environmental effects on the animals' increasing inability
37
38 779 to cope as climate changed. The results in Fig. 4, for example, imply that, historically,
39
40 780 declining populations at high latitudes are likely to have become locked into
41
42 781 mountain-top retreats as climate deteriorated, leading to small, isolated pockets that
43
44 782 are inevitably more vulnerable to extinction in the face of environmental shocks
45
46 783 (Dunbar 1998; Cowlishaw & Dunbar 2000).
47
48
49

50 784 One final puzzle is the fact that gibbons are not found east of the Dieng
51
52 785 Mountains in central Java. Kappeler (1984a,b) attributed this to the kinds of forests
53
54 786 found in the eastern half of the island, which tend to be more deciduous. Fossil
55
56 787 gibbons have been recorded at Trinil, central Java (in the Lower/Middle Pleistocene
57
58
59
60

1
2
3 788 deposits that produced the hominins: Ingicco, Vos & Hoffman 2014) and at the
4
5 789 Ngandong and Gunung Dawung hominin sites in east Java (Storm & Vos 2006) (see
6
7
8 790 also Jablonski & Chapin 2009), so this area clearly did once support gibbon
9
10 791 populations. The fact that west Java was one of the refugia for gibbons during the dry
11
12 792 phases of the late Pleistocene Ice Ages (Brandon-Jones 1998) makes this all the more
13
14 793 puzzling. It may be that anthropogenic factors resulted in their demise, although these
15
16
17 794 are unlikely to have kicked in until historical times.
18
19
20 795
21
22 796

23 797 **Acknowledgements**

24
25
26 798 The authors declare no conflicts of interest. This paper is based entirely on published
27
28 799 data and no ethical issues arise from it. We thank the referees for some helpful
29
30 800 suggestions.
31
32
33 801

REFERENCES

- 802
803
804 Ameca y Juárez, E.I., Mace, G.M., Cowlshaw, G., Cornforth, W.A. & Pettorelli, N.
805 (2013). Assessing exposure to extreme climatic events for terrestrial
806 mammals. *Conservation Letters*, 6, 145–153.
- 807 Alfred, J.R.B. & Sati, J.P. (1986). The gibbon with special reference to *H. hoolock*. In
808 Majupuria, T. C. (Ed.), *Wildlife wealth of India: Resources and*
809 *management*, pp. 384-390. Bangkok: Technical Press Service.
- 810 Alfred, J.R.B. & Sati, J.P. (1990). Survey and census of the hoolock gibbon in west
811 Garo Hills, Northeast India. *Primates*, 31, 299-306.
- 812 Bartlett, T.Q. (2009). Seasonal home range use and defendability in white-handed
813 gibbons (*Hylobates lar*) in Khao Yai National Park, Thailand. In S. Lappan
814 & D.J. Whittaker (Eds.), *The gibbons*, pp. 265-275. New York: Springer.
- 815 Bettridge, C. & Dunbar, R.I.M. (2012). Modelling the biogeography of fossil
816 baboons. *International Journal of Primatology*, 33, 1278-1308.
- 817 Bettridge, C., Lehmann, J. & Dunbar, R. (2010). Trade-offs between time, predation
818 risk and life history, and their implications for biogeography: a systems
819 modelling approach with a primate case study. *Ecological Modelling*, 221,
820 777-790.
- 821 Brandon-Jones, D. (1998). Pre-glacial Bornean primate impoverishment and
822 Wallace's Line. In Hall, R. & Holloway, J.D. (Eds.), *Biogeography and*
823 *geological evolution*, pp. 393-404. Leiden: Backhuys.
- 824 Brockelman, W.Y., Nathalang, A., Greenberg, D.B. & Suwanvecho, U. (2014).
825 Evolution of small-group territoriality in gibbons. In J. Yamagiwa & L.
826 Karczmarski (Eds.), *Primates and cetaceans*, pp. 213-230. Tokyo: Springer.

- 1
2
3 827 Brotherton, P.N.M. & Komers, P.E. (2003). Mate guarding and the evolution of social
4
5 828 monogamy in mammals. In U.H. Reichard & C. Boesch (Eds.), *Monogamy:*
6
7 829 mating strategies and partnerships in birds, humans and other mammals, pp.
8
9 830 42–58. Cambridge: Cambridge University Press.
- 11
12 831 Bryant, J.V., Olson, V.A., Chatterjee, H.J. & Turvey, S.T. (2015). Identifying
13
14 832 environmental versus phylogenetic correlates of behavioural ecology in
15
16 833 gibbons: implications for conservation management of the world’s rarest
17
18 834 ape. *BMC Evolutionary Biology*, 15, 171.
- 20
21 835 Burton-Chellew, M. & Dunbar, R.I.M. (2015). Hamilton’s Rule predicts anticipated
22
23 836 social support in humans. *Behavioral Ecology*, 26, 130-137.
- 24
25 837 Carne, C., Semple, S. & Lehmann, J. (2012). The effects of climate change on
26
27 838 orangutans: a time budget model. *Climate Models*. www.InTechOpen.com.
28
29 839 doi: 10.5772/3324
- 31
32 840 Chapman, C., & Chapman, L. (2000). Determinants of group size in primates: the
33
34 841 importance of travel costs. In S. Boinski & P. A. Garber (Eds.), *On the move:*
35
36 842 how and why animals travel in groups (pp. 24–42). Chicago: University of
37
38 843 Chicago Press.
- 40
41 844 Chivers, D.J. (1984). Feeding and ranging in gibbons: a summary. In Preuschoft,
42
43 845 H.H., Chivers, D.J., Brockelman, W.Y. & Creel, N. (Eds.), *The lesser apes*,
44
45 846 pp. 267-281. Edinburgh: Edinburgh University Press.
- 47
48 847 Clutton-Brock, T. & Janson, C. (2012). Primate socioecology at the crossroads: past,
49
50 848 present, and future. *Evolutionary Anthropology*, 21, 136-150.
- 51
52 849 Coleman, B.T. & Hill, R.A. (2014). Living in a landscape of fear: the impact of
53
54 850 predation, resource availability and habitat structure on primate range
55
56 851 use. *Animal Behaviour*, 88, 165-173.
- 57
58
59
60

- 1
2
3 852 Cowlshaw, G. (1992). Song function in gibbons. *Behaviour*, 121, 131-153.
4
5 853 Cowlshaw, G. (1996). Sexual selection and information content in gibbon song
6
7 bouts. *Ethology*, 102, 272-284.
8
9 855 Cowlshaw, G. & Dunbar, R.I.M. (2000). Primate conservation biology. Chicago:
10
11 Chicago University Press.
12
13 856
14 857 Dàvid-Barrett, T. & Dunbar, R.I.M. (2016). Bipedality and hair loss in human
15
16 evolution revisited: the impact of altitude and activity scheduling. *Journal of*
17
18 *Human Evolution*, 94, 72-82.
19
20 859
21 860 Dunbar, R.I.M. (1980). Determinants and evolutionary consequences of dominance
22
23 among female gelada baboons. *Behavioural Ecology and Sociobiology*, 7, 253-265.
24
25 861
26 862 Dunbar, R.I.M. (1992a). Time: a hidden constraint on the behavioural ecology of
27
28 baboons. *Behavioural Ecology and Sociobiology*, 31, 35-49.
29
30 863
31 864 Dunbar, R.I.M. (1992b). A model of the gelada socioecological system. *Primates*, 33,
32
33 69-84.
34
35 866 Dunbar, R.I.M. (1995a). The mating system of Callitrichid primates. I. Conditions for
36
37 the coevolution of pairbonding and twinning. *Animal Behaviour* 50: 1057-
38
39 1070.
40
41 868
42 869 Dunbar, R.I.M. (1995b) The mating system of Callitrichid primates. II. The impact of
43
44 helpers. *Animal Behaviour* 50: 1071-1089.
45
46 871 Dunbar, R.I.M. (1998). Impact of global warming on the distribution and survival of
47
48 the gelada baboon: a modelling approach. *Global Change Biology*, 4, 293-304.
49
50 872
51 873 Dunbar, R.I.M. (2002). Modelling primate behavioural ecology. *International Journal of*
52
53 *Primatology*, 23, 785-819.
54
55
56
57
58
59
60

- 1
2
3 875 Dunbar, R.I.M. (2018a). Social structure as a strategy to mitigate the costs of group-
4
5 876 living: a comparison of gelada and guereza monkeys. *Animal Behaviour*, 136,
6
7 877 53-64.
8
9
10 878 Dunbar, R.I.M. (2018b). The anatomy of friendship. *Trends in Cognitive Science*, 22,
11
12 879 32-51.
13
14 880 Dunbar, R.I.M. & Lehmann, J. (2013). Grooming and social cohesion in primates: a
15
16 881 comment on Grueter et al. *Evolution and Human Behaviour*, 34, 453-455.
17
18 882 Dunbar, R.I.M. & Shi, J. (2013). Time as a constraint on the distribution of feral goats
19
20 883 at high latitudes. *Oikos*, 122, 403-410.
21
22 884 Dunbar, R.I.M. & Shultz, S. (2010). Bondedness and sociality. *Behaviour*, 147, 775-803.
23
24 885 Dunbar, R., Korstjens, A. & Lehmann, J. (2009). Time as an ecological constraint.
25
26 886 *Biological Reviews*, 84, 413-429.
27
28 887 Dunbar, R., MacCarron, P. & Shultz, S. (2018a). Primate social group sizes exhibit a
29
30 888 regular scaling pattern with natural attractors. *Biology Letters*, 14, 20170490.
31
32 889 Dunbar, R.I.M., MacCarron, P. & Robertson, C. (2018b). Tradeoff between fertility
33
34 890 and predation risk drives a geometric sequence in the pattern of group sizes in
35
36 891 baboons. *Biology Letters*, 14, 20170700
37
38 892 Ellefson, J.O. (1974). A natural history of the white-handed gibbon in the Malayan
39
40 893 Peninsula. In Rumbaugh, D.M. (Eds.), *Gibbon and siamang*, Vol. 3, pp. 1-
41
42 894 136. Basel: Karger.
43
44 895 Fan, P. (2017). The past, present, and future of gibbons in China. *Biological*
45
46 896 *Conservation*, 210, 29-39..
47
48 897 Fei, H.-L., Zhang, D., Yuan, S.D., Zhang, L. & Fan, P-F. (2017). Antipredation
49
50 898 sleeping behavior of skywalker hoolock gibbons (*Hoolock tianxing*) in Mt.
51
52 899 Gaoligong, Yunnan, China. *International Journal of Primatology*, 38, 629-
53
54
55
56
57
58
59
60

- 1
2
3 900 641.
4
5 901 Gallagher, A.J., Creel, S., Wilson, R.P. & Cooke, S.J. (2017). Energy landscapes and
6
7 902 the landscape of fear. *Trends in Ecology & Evolution*, 32, 88-96.
8
9
10 903 Heldstab, S.A., Schaik, C.P. & Isler, K. (2017). Getting fat or getting help? How
11
12 904 female mammals cope with energetic constraints on reproduction. *Frontiers in*
13
14 905 *Zoology*, 14, 29.
15
16
17 906 Hill, R.A., Lycett, J.E. & Dunbar, R.I.M. (2000). Ecological determinants of birth
18
19 907 intervals in baboons. *Behavioral Ecology*, 11, 560-564.
20
21
22 908 Hill, R.A. (1999). Ecological and demographic determinants of time budgets in
23
24 909 baboons: implications for cross-population models of baboon socioecology.
25
26 910 PhD thesis, University of Liverpool.
27
28
29 911 Ingicco, T., de Vos, J. & Huffman, O.F. (2014). The oldest gibbon fossil
30
31 912 (Hylobatidae) from insular Southeast Asia: evidence from Trinil (East Java,
32
33 913 Indonesia), Lower/Middle Pleistocene. *PLoS ONE*, 9, e99531.
34
35
36 914 Jablonski, N.G. & Chaplin, G. (2009). The fossil record of gibbons. In S. Lapan & D.
37
38 915 Whittaker (Eds.), *The gibbons*, pp. 111-130. New York: Springer.
39
40 916 Janson, C. H., & van Schaik, C. P. (1988). Recognizing the many faces of primate
41
42 917 food competition: methods. *Behaviour*, 105, 165–186.
43
44
45 918 Johns, A. (1986). Effects of selective logging on the behavioural ecology of west
46
47 919 Malaysian primates. *Ecology*, 67, 684-694.
48
49
50 920 Kamilar, J.M. & Cooper, N. (2013). Phylogenetic signal in primate behaviour,
51
52 921 ecology and life history. *Philosophical Transactions of the Royal Society,*
53
54 922 *London*, 368B, 20120341.
55
56
57 923 Kappler, M. (1984a). The gibbon in Java. In Preuschoft, H.H., Chivers, D.J.,
58
59 924 Brockelman, W.Y. & Creel, N. (Eds.), *The lesser apes*, pp. 19-31. Edinburgh:
60

- 1
2
3 925 Edinburgh University Press.
4
5
6 926 Kappeler, M. (1984b). Diet and feeding behaviour of the moloch gibbon. In
7
8 927 Preuschoft, H.H., Chivers, D.J., Brockelman, W.Y. & Creel, N. (Eds.), The
9
10 928 lesser pes, pp. 228-241. Edinburgh: Edinburgh University Press.
11
12 929 Komers, P.E. & Brotherton, P.N.M. (1997). Female space use is the best predictor of
13
14 930 monogamy in mammals. *Proceedings of the Royal Society, London*, 264B,
15
16 931 1261–1270.
17
18
19 932 Korstjens, A. & Dunbar, R. (2007). Time constraints limit group sizes and distribution
20
21 933 in red and black-and-white colobus monkeys. *International Journal of*
22
23 934 *Primatology*, 28, 551-575.
24
25
26 935 Korstjens, A., Lehmann, J. & Dunbar, R. (2010). Resting time as an ecological
27
28 936 constraint on primate biogeography. *Animal Behaviour*, 79, 361-374.
29
30
31 937 Korstjens, A.H., Lehmann, J. & Dunbar, R.I.M. (2018). Time constraints do not limit
32
33 938 group size in arboreal guenons but do explain community size and distribution
34
35 939 patterns. *International Journal of Primatology* (in press).
36
37
38 940 Korstjens, A., Verhoeckx, I. & Dunbar, R. (2006). Time as a constraint on group size in
39
40 941 spider monkey. *Behavioural Ecology and Sociobiology*, 60, 683-694.
41
42
43 942 Kudo, H. & Dunbar, R.I.M. (2001). Neocortex size and social network size in primates.
44
45 943 *Animal Behaviour*, 62, 711-722.
46
47 944 Laundré, J.W., Hernández, L., Medina, P.L., Campanella, A., López-Portillo, J.,
48
49 945 González-Romero, A., Grajales-Tam, K.M., Burke, A.M., Gronemeyer, P. &
50
51 946 Browning, D.M. (2014). The landscape of fear: the missing link to understand
52
53 947 top-down and bottom-up controls of prey abundance? *Ecology*, 95, 1141-1152.
54
55
56 948 Lehmann, J., Korstjens, A. & Dunbar, R.I.M. (2007a). Group size, grooming and
57
58 949 social cohesion in primates. *Animal Behaviour*, 74, 1617-1629.
59
60

- 1
2
3 950 Lehmann, J., Korstjens, A. & Dunbar, R.I.M. (2007b). Fission-fusion social systems
4
5 951 as a strategy for coping with ecological constraints: a primate case.
6
7 952 *Evolutionary Ecology*, 21, 613-634.
8
9
10 953 Lehmann, J., Korstjens, A. & Dunbar, R.I.M. (2008a). Time management in great apes:
11
12 954 implications for gorilla biogeography. *Evolutionary Ecology Research*, 10,
13
14 955 515-536.
15
16
17 956 Lehmann, J., Korstjens, A. & Dunbar, R.I.M. (2008b). Time and distribution: a model
18
19 957 of ape biogeography. *Ethology, Ecology and Evolution*, 20, 337-359.
20
21
22 958 Lehmann, J., Korstjens, A. & Dunbar, R.I.M. (2010). Apes in a changing world – the
23
24 959 effects of global warming on the behaviour and distribution of African apes.
25
26 960 *Journal of Biogeography*, 37, 2217-2231.
27
28
29 961 Le Houérou, H.N. & Hoste, C.H. (1977). Rangeland production and annual rainfall
30
31 962 relations in the Mediterranean Basin and in the African Sahelo-Sudanian
32
33 963 zone. *Journal of Range Management*, 30, 181-189.
34
35
36 964 Liu, K. Shen, C. & Louie, K. (2001). A 1,000-year history of typhoon landfalls in
37
38 965 Guangdong, Southern China, reconstructed from Chinese historical
39
40 966 documentary records. *Annals of the Association of American Geographers*, 91,
41
42 967 453–464.
43
44
45 968 Lo Seen Chong, D., Mougin, E. & Gastellu-Etchegorry, J.P. (1993). Relating the
46
47 969 global vegetation index to net primary productivity and actual
48
49 970 evapotranspiration over Africa. *International Journal of Remote Sensing*, 14,
50
51 971 1517-1546.
52
53
54 972 Lukas, D. & Clutton-Brock, T.H. (2013). The evolution of social monogamy in
55
56 973 mammals. *Science*, 341, 526–530.
57
58
59 974 Marshall, A.J. & Leighton, M. (2006). How does food availability limit the population
60

- 1
2
3 975 density of white-bearded gibbons? In Hohmann, G., Robbins, M.M. & Boesch, C.
4
5 976 (Eds.), *Feeding ecology in apes and other primates*, pp. 311-333. Cambridge:
6
7 Cambridge University Press.
8
9
10 978 Marshall, H. H., Carter, A. J., Rowcliffe, J. M., & Cowlshaw, G. (2012). Linking
11
12 979 social foraging behaviour with individual time budgets and emergent group-level
13
14 980 phenomena. *Animal Behaviour*, 84, 1295–1305.
15
16
17 981 Mount, L.E. (1979). *Adaptation to thermal environment*. London: Arnold.
18
19 982 Opie, K., Atkinson, Q., Dunbar, R. & Shultz, S: Infanticide still drives primate
20
21 983 monogamy. *Proceedings of the National Academy of Sciences, USA*, 111,
22
23 984 E1675.
24
25
26 985 Palombit, R. (1995). Inter- and intraspecific variation in the diets of sympatric
27
28 986 siamang (*Hylobates syndactylus*) and lar gibbons (*Hylobates lar*). *Folia*
29
30 987 *Primatologica*, 68, 321-337.
31
32
33 988 Palombit, R.A. (1997). Inter- and intraspecific variation in the diets of sympatric
34
35 989 siamang (*Hylobates syndactylus*) and lar gibbons (*Hylobates lar*). *Folia*
36
37 990 *Primatologica*, 68, 321-337.
38
39
40 991 Peters, R.H. (1983). *Ecological implications of body size*. Cambridge: Cambridge
41
42 992 University Press.
43
44
45 993 Pollard, K. A., & Blumstein, D. T. (2008). Time allocation and the evolution of group
46
47 994 size. *Animal Behaviour*, 76, 1683–1699.
48
49 995 Raemaekers, J.J. (1979). Ecology of sympatric gibbons. *Folia Primatologica*, 31, 227-
50
51 996 245.
52
53
54 997 Reichard, U.H. (2003). Social monogamy in gibbons: the male perspective. In U.H.
55
56 998 Reichard & C. Boesch (Eds.), *Monogamy: mating strategies and partnerships*
57
58 999 *in birds, humans and other mammals*, pp. 190-213. Cambridge: Cambridge
59
60

- 1
2
3 1000 University Press.
4
5 1001 Reiter, P. (2000). From Shakespeare to Defoe: malaria in England in the Little Ice
6
7 1002 Age. *Emerging Infectious Diseases*, 6, 1–11.
8
9
10 1003 Riginos, C., 2015. Climate and the landscape of fear in an African savanna. *Journal of*
11
12 1004 *Animal Ecology*, 84, 124-133.
13
14 1005 Rosenzweig, M.L. (1968). Net primary productivity of terrestrial communities:
15
16 1006 prediction from climatological data. *American Naturalist*, 102, 67-74.
17
18 1007 Rutberg, A.T. (1983). The evolution of monogamy in primates. *Journal of Theoretical*
19
20 1008 *Biology*, 104, 93-112.
21
22 1009 van Schaik, C.P. (1983). Why are diurnal primates living in groups? *Behaviour*, 87,
23
24 1010 120-144.
25
26
27 1011 van Schaik, C.P. & Dunbar, R.I.M. (1990). The evolution of monogamy in large-
28
29 1012 bodied primates: a new hypothesis and some critical tests. *Behaviour*, 115, 30-
30
31 1013 62.
32
33
34 1014 van Schaik, C.P. & van Hooff, J. (1983). On the ultimate causes of primate social
35
36 1015 systems. *Behaviour*, 85, 91-117.
37
38
39 1016 van Soest, P.J. (1982). Nutritional ecology of the ruminant. Ithaca, NY: Cornell
40
41 1017 University Press.
42
43
44 1018 Seyfarth, R.M. & Cheney, D.L. (1984). Grooming, alliances and reciprocal altruism in
45
46 1019 vervet monkeys. *Nature*, 308, 541.
47
48
49 1020 Shultz, S., Noe, R., McGraw, S. & Dunbar, R.: A community-level evaluation of the
50
51 1021 impact of prey behavioural and ecological characteristics on predator diet
52
53 1022 composition. *Proc. R. Soc. Lond.*, 271B, 725-732.
54
55
56
57
58
59
60

- 1
2
3 1023 Shultz, S. & Finlayson, L.V. (2010). Large body and small brain and group sizes are
4
5 1024 associated with predator preferences for mammalian prey. *Behavioral*
6
7 1025 *Ecology*, 21, 1073-1079.
- 8
9
10 1026 Shultz, S., Noe, R., McGraw, S. & Dunbar, R.I.M. (2004). A community-level
11
12 1027 evaluation of the impact of prey behavioural and ecological characteristics on
13
14 1028 predator diet composition. *Proceedings of the Royal Society, London*, 271B, 725-
15
16 1029 732.
- 17
18
19 1030 Snaith, T.V. & Chapman, C.A. (2007). Primate group size and interpreting
20
21 1031 socioecological models: do folivores really play by different
22
23 1032 rules?. *Evolutionary Anthropology*, 16, 94-106.
- 24
25
26 1033 Srikosamatara, S. (1984). Ecology of pileated gibbons in southeast Thailand. In
27
28 1034 Preuschoft, H.H., Chivers, D.J., Brockelman, W.Y. & Creel, N. (Eds.), *The*
29
30 1035 *lesser apes*, pp. 242-257. Edinburgh: Edinburgh University Press.
- 31
32
33 1036 Storm, P. & de Vos, J. (2006). Rediscovery of the Late Pleistocene Punung hominin
34
35 1037 sites and the discovery of a new site Gunung Dawung in East
36
37 1038 Java. *Palaeobiodiversity and Palaeoenvironments*, 86, 271-281.
- 38
39
40 1039 Sutcliffe, A.J., Dunbar, R.I.M., Binder, J. & Arrow, H. (2012). Relationships and the
41
42 1040 social brain: integrating psychological and evolutionary perspectives. *British*
43
44 1041 *Journal of Psychology*, 103, 149-168.
- 45
46
47 1042 Tolon, V., Dray, S., Loison, A., Zeileis, A., Fischer, C. & Baubet, E. (2009).
48
49 1043 Responding to spatial and temporal variations in predation risk: space use of a
50
51 1044 game species in a changing landscape of fear. *Canadian Journal of*
52
53 1045 *Zoology*, 87, 1129-1137.
- 54
55
56 1046 Turvey, S.T., Crees, J.J. & Di Fonzo, M.M.I. (2015). Historical data as a baseline for
57
58 1047 conservation: reconstructing long-term faunal extinction dynamics in Late
59
60

- 1
2
3 1048 Imperial–modern China. *Proceedings of the Royal Society, London, B*
4
5 1049 20151299.
6
7
8 1050 Willems, E.P. & Hill, R.A. (2009). A critical assessment of two species distribution
9
10 1051 models: a case study of the vervet monkey (*Cercopithecus aethiops*). *Journal*
11
12 1052 *of Biogeography*, 36, 2300-2312.
13
14 1053 Williamson, D.K. & Dunbar, R.I.M. (1998). Energetics, time budgets and group size.
15
16 In Lee, P.C. (Ed.), *Comparative primate Ssocioecology*, pp. 320-338.
17 1054
18 Cambridge: Cambridge University Press.
19 1055
20
21 1056 Wittenberger, J.F. & Tilson, R.L. (1980). The evolution of monogamy: hypotheses
22
23 and evidence. *Annual Review of Ecology and Systematics*, 11, 197-232.
24 1057
25
26 1058 Wrangham, R.W., Gittleman, J.L. & Chapman, C.A. (1993). Constraints on group
27
28 size in primates and carnivores: population density and day-range as assays
29 1059
30 of exploitation competition. *Behavioral Ecology and Sociobiology*, 32, 199-
31 1060
32 209.
33 1061
34
35 1062
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Legends to Figures

1064

1065

1066 Fig. 1. Flow chart for the gibbon time budget model, summarising the causal
1067 relationships between climatic and behavioural variables based on the
1068 equations given in Table 1. Solid lines: positive effects; dashed lines: negative
1069 effects.

1070

1071 Fig. 2. Time spent moving plotted against mean population group size. Dashed line is
1072 least squares regression line. Filled symbols: gibbons; unfilled symbols:
1073 siamang. Source: online *Dataset S1*.

1074

1075 Fig. 3. (a) Maximum ecologically tolerable group size predicted by the time budget
1076 model, plotted against observed mean group size for different gibbon (filled
1077 symbols) and siamang (unfilled symbols) populations. (b) Predicted maximum
1078 group size allowing for a 15% increase in the length of the active day. The solid
1079 line demarcates the minimum group size for demographic viability (2 adults plus
1080 0.96 immatures: see text for details); dashed line in (a) is the line of equilibrium
1081 (N_{max} = observed mean).

1082

1083 Fig. 4. Maximum group size predicted by the time budget model for different
1084 combinations of (a) latitude and altitude and (b) annual rainfall and mean
1085 temperature.

1086

1087 Fig. 5. Mean ($\pm 95\%$ CI) maximum group size (N_{max}) predicted by the time budget
1088 model for habitats within different geographic regions. Filled circles: regions

1
2
3 1089 where gibbons currently live; unfilled circles: regions where gibbons do not
4
5 1090 currently live; grey circle: gibbons historically present before 1800 AD. Dashed
6
7 1091 line: minimum group size for demographic viability (2.96 individuals: see text).
8
9
10 1092 Present: 18 sites within the current biogeographic distribution of gibbons within
11
12 1093 Indo-China, Sumatra and Borneo. W Java: 3 sites in west Java where gibbons
13
14 1094 occur; E Java: 8 sites in Java east of the Dieng Mountains where gibbons do not
15
16 1095 currently occur; E Indonesia: 2 sites on Celebes and western Papua New Guinea,
17
18 1096 east of the Wallace Line where gibbons do not occur; NW and NE India: 8 and 7
19
20 1097 sites, respectively, in northwest and northeast India (including Bangladesh) where
21
22 1098 gibbons do not occur; S India: 7 sites in southern India plus Sri Lanka; SW
23
24 1099 China: 3 sites within current gibbon range in Yunnan and Hainan. SE China: one
25
26 1100 site from each province in southern China where gibbons were recorded as being
27
28 1101 present in historical documents before 1800 AD (N=14 site). N China: one site
29
30 1102 from each province in northern China (including Tibet) where gibbons were not
31
32 1103 recorded historically (N=7 sites). Locations and climate data from online *Dataset*
33
34 1104 *S2*.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1106 Fig. 6. Mean ($\pm 95\%$ CI) (a) feeding time, (b) moving time and (c) enforced rest time
1107 predicted by the time budget model for the regional sites shown in Fig. 5. Moving
1108 time does not include an adjustment for group size. Regions and symbols as for
1109 Fig. 5.

1110
1111 Fig. 7. (a) Mean ($\pm 95\%$ CI) ratio of observed to predicted time spent feeding and
1112 resting for populations for which predicted $N_{max} < 5$ (unfilled symbols) or N_{max}
1113 > 5 (filled symbols). Predicted rest time is enforced rest time. (b) Difference in

1
2
3 1114 rest time (observed minus predicted by model) for individual hylobatid
4
5 1115 populations plotted against difference in group size (observed minus *Nmax*).
6

7
8 1116 Source: Dataset S1.
9

10 1117
11

12 1118 Fig. 8. Mean observed group size for individual populations plotted against absolute
13

14 1119 latitude for the population. The best fit equation (dashed line) is Group size =
15

16 1120 $3.71 - 0.014*Lat + 0.012*Lat^2 - 0.0004*Lat^3$ ($r^2=0.159$, $F_{3,50}=3.15$, $p=0.033$).
17
18

19 1121 Source: Dataset S1.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Regression equations for the hylobatid model.

Variable	Equation	r ²	F	df	p
(1) Feed time (%)	$41.634 - 0.003 * \text{RAIN} - 0.009 * \text{ALT} + 0.363 * \text{LAT}_{\text{abs}}^{\Omega}$	0.501	5.03	3,15	0.013
(2) Moving time (%)	$17.316 + 0.462 * \text{LAT}_{\text{abs}}$	0.166	3.59	1,18	0.074
(2a) Moving time (%)	$5.297 + 0.290 * \text{LAT}_{\text{abs}} + 3.080 * \text{N}$	0.220	1.98	2,14	0.175
(3) Rest _{ENF} time (%)¶	$-23.24 + 1.33 * \text{TEMP} + 0.259 * \text{LEAF} + 6.12 * \text{T}_{\text{moSD}}$	from Korstjens et al. (2010)			
(4) Social(%)†	$2.968 + 0.109\text{N}$	0.440	29.90	1,38	<0.0001
(5) Time budget	$100 = \text{FEED} + \text{MOVE} + \text{REST}_{\text{ENF}} + \text{SING} + \text{SOCIAL}$				
(5a)	$100 = \text{FEED} + (5.297 + 0.290 * \text{LAT}_{\text{abs}} + 3.080 * \text{N}) + (\text{REST}_{\text{ENF}} + \text{REST}_{\text{FREE}}) + \text{SING} + (2.968 + 0.109 * \text{N})$				
(5b) Nmax§	$= ((100 - \text{FEED} - (5.297 + 0.290 * \text{LAT}_{\text{abs}}) - \text{REST} - 4.96^{\ddagger}) + 2.968) / (0.109 + 3.080)$	by substitution			

Ω Absolute latitude

¶ Rest time consists of two components: enforced rest (REST_{ENF}, imposed on the animal by climatic conditions and its dietary strategy) and free rest (REST_{FREE}) that represents uncommitted time that can be allocated to any other activity when required (see Korstjens et al. 2010).

† Recalculated from Lehmann et al (2007a)

§ Maximum ecologically tolerable group size (the maximum group size that will allow the time budget to be balanced)

‡ 4.96 = average percentage of day devoted to singing by gibbons

Table 2. Supplementary equations

Variable	Equation	r ²	F	df	p
TmoSD **	10.440 + 0.093*LAT _{Abs} - 0.001*ALT - 0.821*TEMP + 0.588*TEMP _{min}	0.597	17.75	4,48	<0.0001
Mean temperature (°C)	27.651 - 0.003*ALT - 0.158*LAT _{Abs}	0.672	60.33	2,59	0.0001
Altitude (m)	4145.354 - 147.236*TEMP + 0.023*RAIN	0.575	37.14	2,55	<0.0001
Temp _{var} †	10.196				
Absolute Latitude	48.732 - 0.004*RAIN - 1.142*TEMP	0.580	35.23	2,51	<0.0001
Rain	3279.523 - 53.609*LAT _{Abs} + 0.025*ALT	0.286	10.60	2,52	<0.0001
Temp _{min} (°C)	-5.776 + 1.033*TEMP	0.922	623.63	1,53	<0.0001
Territory size (ha)	2498.88 + 0.077*ALT - 477.830*TEMP _{min} + 401.718*TEMP _{max} - 477.03*TEMP _{var} - 32.260*TEMP _{msoSD} - 0.032*RAIN	0.894	19.60	6,14	<0.0001
Leaf in diet (%)	99.430 - 1.000*LAT _{abs} + 0.009*ALT + 7.012*TEMP _{var} + 7.879*TEMP _{min} - 9.779*TEMP _{max}	0.611	5.04	5,16	0.006
Fruit in diet (%)	-51.27 + 10.65*TEMP - 7.71*TEMP _{min}	0.382	6.19	2,20	0.008
Active day (hr) ^Ω	11.273 - 0.001*RAIN	0.905	19.15	1,2	0.048

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

** Standard deviation of monthly mean temperatures across the year

† Difference between mean monthly maximum and mean monthly minimum temperature

Ω Gibbons only

For Peer Review

Table 3. Time budget differences between siamang and gibbons

Variable	Siamang		Gibbons*	
	Mean±SD	N§	Mean±SD	N§
% fruit in diet	40.7±16.1	6	57.9±18.7	22
% leaf in diet	34.6±11.6	4	33.7±30.4	22
Feed (%)	49.6±6.8	5	34.4±8.5	19
Move (%)	18.0±5.8	5	21.4±7.9	16
Rest (%)	34.4±10.1	5	34.7±9.8	18
Sing (%)	2.0	1	4.96±3.1	10
Social time (%)	3.0	1	5.87±3.5	12
Active day (hr)	10.1±0.3	2	9.06±0.9	7
Net difference in available time†		16.8%		

* sites used in time budget analyses only

§ Number of studies

† Equivalent change in gross time budget achieved by the siamang compared to gibbons (~3% by reducing singing, ~3% by reducing social time and ~11% by increasing length of active day)

Table 4. Comparison of climatic variables between southwest China (where gibbons currently exist) and southeast China (where they were present historically, but no longer are).

Provinces of China	Gibbons present	TEMP	TEMP _{min}	TEMP _{max}	TEMP _{var}	TEMP _{moSD}	RAIN
Southwest	current	20.6	16.5	24.6	8.1	3.8	1973
Southeast	historical	16.6	12.5	20.9	9.7	8.1	1163
	F _{1,15}	2.50	1.88	2.61	0.21	19.97	9.22
	p	0.135	0.191	0.127	0.657	0.0005	0.008

Environment and Time as Constraints on the Biogeographical Distribution of Gibbons

R.I.M. Dunbar, Susan M. Cheyne, Daoying Lan, Amanda Korstjens,
Julia Lehmann & G. Cowlshaw

Online Supplementary Material

Alternative model equations

Table S1 provides alternative multivariate equations for feeding and moving time generated by the backwards regression model. Most of the slope coefficients are similar across the alternative equations. However, note that the magnitude of the effect of group size on moving time for Equation (2c) is considerably less. Nonetheless, the values for both Equations (2a) and (2c) straddle the observed value for the chimpanzee model.

Table S2 gives the correlation values for maximum ecologically tolerable group size predicted by the original model based on Table 1 in the main text and the various alternative feeding and moving time equations given in Table S1. All alternative statistical models yield predicted maximum group sizes that correlate significantly with those predicted by the original model based on the equations of Table 1.

Fig. S1 plots predicted maximum group size for individual study sites in the main dataset (online *Dataset S1*) against the equivalent value predicted by substituting the various alternative feeding and moving time equations. Alternative feeding time equations do not have a significant effect on predicted maximum group size. Note that the alternative moving time equation predicts slightly larger group sizes than those predicted by the original model of Table 1 because the effect of group size on moving time in Equation 2c is lower (1.72 vs 3.08), yielding slightly larger group sizes. Nonetheless, the difference in the size of the largest groups predicted is modest (25 vs 15), the outcomes do not differ significantly (Table S2). Hence, using these alternative equations does not change any of the main conclusions.

Table S1. Alternative regression equations for feeding and moving time for the hylobatid model.

Variable	Equation	r^2	F	df	p
(1) Feed time (%)	$41.634 + 0.363 * LAT_{abs} - 0.009 * ALT - 0.003 * RAIN$	0.501	5.03	3,15	0.013
(1a)	$53.955 + 0.243 * LAT - 0.010 * ALT - 0.472 * TEMP + 0.243 * TEMPVAR - 0.004 * RAIN$	0.340	2.86	5,13	0.059
(1b)	$55.879 + 0.279 * LAT_{abs} - 0.010 * ALT - 0.485 * TEMP - 0.004 * RAIN$	0.520	3.80	4,14	0.027
(1c)	$33.230 + 0.469 * LAT_{abs} - 0.009 * ALT$	0.461	6.85	2,16	0.007
(2) Moving time (%)	$17.316 + 0.462 * LAT_{abs}$	0.273	5.25	1,14	0.038
(2a)	$5.297 + 0.290 * LAT_{abs} + 3.080 * N$	0.220	1.98	2,14	0.175
(2b)	$29.196 + 0.306 * LAT - 0.365 * TEMP_{max}$	0.294	2.71	2,13	0.104
(2c)	$19.409 + 0.303 * LAT_{abs} - 0.275 * TEMP_{max} + 1.718 * N$	0.312	1.81	3,12	0.199

Table S2. Correlations between maximum group size predicted by the original model of Table 1 and a model using the different equations for feeding and moving time given in Table S1, for the actual hylobatid study sites (as listed in online *Dataset S1*). In each case, one equation is substituted, with all the remaining equations as in the original model of Table 1.



Pearson Correlations

		Feed Eqn 1a	Feed Eqn 1b	Feed Eqn 1c	Move Eqn 2c	Feed Eqn 1a with Move Eqn 2c	Feed Eqn 1c with Move Eqn 2c
Original model	Correlation	0.994	0.995	0.988	0.996	0.987	0.986
	p (2-tailed)	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
	N	51	51	51	51	51	51
Feed Eqn 1a	Correlation		0.999	0.974	0.996	0.997	0.978
	p (2-tailed)		<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
	N		51	51	51	51	51
Feed Eqn 1b	Correlation			0.975	0.997	0.997	0.980
	p (2-tailed)			<0.0001	<0.0001	<0.0001	<0.0001
	N			51	51	51	51
Feed Eqn 1c	Correlation				0.984	0.966	0.997
	p (2-tailed)				<0.0001	<0.0001	<0.0001
	N				51	51	57

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Move Eqn 2c	Correlation					0.995	0.989
	p (2-tailed)					<0.0001	<0.0001
	N					51	51
Feed Eqn 1a with Move Eqn 2c	Correlation						0.977
	p (2-tailed)						<0.0001
	N						51

For Peer Review

Figure S1

Maximum group size predicted by the original model using the Table 1 equations plotted against maximum group size for all populations in the sample of studies (*Dataset SI*), using the alternative Feed and Move equations given in Table S1. Except for the last pair of graphs, each alternative equation is substituted individually, and all other equations are held constant as in the original model of Table 1. Filled symbols: gibbon populations; unfilled symbols: siamang populations. Dashed line: line of equivalence (the predicted maximum group size is the same for both models).






Fig. S1 (contd)



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Fig. S1 (contd)




Model predictions

Fig. S2 plots the mean ($\pm 95\%$ CI) predicted maximum group size (N_{max}) for sites in countries where hylobatids are known to live and those in adjacent geographical regions (which should be accessible to them) where they do not. These values are calculated from the climate data in *Dataset S2* for those regions listed as within their current geographic range, plus SE China (essentially the provinces where gibbons occurred prior to 1800), NE India (the area north and west of the Brahmaputra in Bangladesh and northeastern India), and central and eastern Java. Indonesia east of the Wallace Line is excluded.

The difference is highly significant (means 2.92 vs 6.8, $N=31$ and 23, respectively; $F_{1,52}=10.71$, $p=0.002$). Note that, in areas where gibbons do not live, half the N_{max} values are below the group size required for demographic viability (2.96 individuals), and most are below the observed mean size of 4.01 individuals. In contrast, all values for locations where gibbons live are above the observed mean group size.

Figure S2

Mean ($\pm 95\%$ CI) predicted maximum group size (N_{max}) for sites in countries where hylobatids are currently found, compared to that for geographically adjacent areas where they do not now occur (see main text for details). The solid line is the calculated minimum group size for demographic viability (2.96 individuals of all ages: see main text); the dashed line is the observed mean group size across all gibbon populations. Source: *Dataset S2*



Speed of travel

We calculate average speed of travel across the day as day journey length (in metres) divided by the number of hours in the day (12) multiplied by the proportion of the day devoted to moving. Speed of travel is weakly predicted by latitude, altitude and temperature:


$$\text{SPEED (m/hr)} = -621.2 - 25.7 * \text{LAT}_{\text{Abs}} + 0.4 * \text{ALT} + 46.5 * \text{TEMP}$$

($r^2=0.674$, $F_{3,7}=1.94$, $p=0.211$).

Fig. S3(a) plots speed against absolute latitude and mean temperature. There is a very slight suggestion that the relationship might be quadratic (U-shaped) in temperature, with speed increasing when ambient temperatures are both low and high, with a minimum at temperatures around 20°C. Alternatively, the relationship may be better described by a power curve in which speed increases dramatically above ~22°C at latitudes <5° either side of the equator.

Figure S3(a)


Mean speed for individual gibbon (filled symbols) and siamang (unfilled symbols) plotted against mean temperature and absolute latitude of the study site, assuming a 12-hour tropical day.




1
2
3
4 Many gibbon populations retire to nest several hours before sunset, with length of
5 active daylength best predicted by mean rainfall (Table 2). To check whether a shorter
6 active day makes any difference to these results, we recalculated speed as a function
7 of actual length of active day using the equation given in Table 2 to predict length of
8 active day from rainfall for each site. Aside from a slight uniform increase in speed,
9 the transformation has no effect (Fig. S3b).
10
11
12
13
14

15 **Figure S3(b)**

16 Mean speed for individual gibbon (filled symbols) and siamang (unfilled symbols) plotted against
17 mean temperature and absolute latitude of the study site, with speed calculated for active day length
18 (estimated using the equation in rainfall from Table 2).
19
20
21
22





Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig. 2

165x131mm (300 x 300 DPI)




FIG. 3B

165x131mm (300 x 300 DPI)




Fig. 4A

123x123mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig. 4B

123x123mm (300 x 300 DPI)




FIG. 5

165x131mm (300 x 300 DPI)




Fig. 6A

165x131mm (300 x 300 DPI)




Fig. 6B

165x131mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig. 6C

165x131mm (300 x 300 DPI)




Fig. 7A

165x131mm (300 x 300 DPI)




Fig. 7B

165x131mm (300 x 300 DPI)

